

UNITED STATES DEPARTMENT OF THE INTERIOR  
BUREAU OF OCEAN ENERGY MANAGEMENT  
GULF OF MEXICO OCS REGION  
NEW ORLEANS, LOUISIANA

SITE-SPECIFIC ENVIRONMENTAL ASSESSMENT

OF

EXPLORATION PLAN  
NO. N-10117

FOR

ANADARKO PETROLEUM CORPORATION

October 2, 2020

**RELATED ENVIRONMENTAL DOCUMENT**

Gulf of Mexico OCS Oil and Gas Lease Sales: 2017-2022  
Gulf of Mexico Lease Sales 249, 250, 251, 252, 253, 254, 256, 257, 259, and 261  
Final Environmental Impact Statement  
(OCS EIS/EA BOEM 2017-009)

Gulf of Mexico OCS Lease Sale  
Final Supplemental Environmental Impact Statement 2018  
(OCS EIS/EA BOEM 2017-074)

Biological Opinion Oil and Gas Leasing, Exploration, Development, Production, Decommissioning,  
and All Related Activities in the Gulf of Mexico Outer Continental Shelf  
(FWS April 20, 2018)

Biological Opinion of the Federally Regulated Oil and Gas Program Activities in the Gulf of Mexico  
(NMFS March 13, 2020)

## FINDING OF NO SIGNIFICANT IMPACT (FONSI)

The Bureau of Ocean Energy Management (BOEM) has prepared a Site-Specific Environmental Assessment (SEA) (No. N-10117) complying with the National Environmental Policy Act (NEPA). NEPA regulations under the Council on Environmental Quality (CEQ) (40 CFR § 1501.3 and § 1508.9), the Department of the Interior (DOI) NEPA implementing regulations (43 CFR § 46), and BOEM policy require an evaluation of proposed major federal actions, which under BOEM jurisdiction includes approving a plan for oil and gas exploration or development activity on the Outer Continental Shelf (OCS).

NEPA regulation 40 CFR § 1508.27(b) requires significance to be evaluated in terms of context and intensity. The context and intensity of impacts caused by similar actions to that proposed were examined at a basin-wide scale in the Gulf of Mexico (GOM) in the following NEPA documents:

- Gulf of Mexico OCS Oil and Gas Lease Sales: 2017-2022 Gulf of Mexico Lease Sales 249, 250, 251, 252, 253, 254, 256, 257, 259, and 261-Final Environmental Impact Statement (Multisale EIS) (OCS EIS/EA BOEM 2017-009);
- Gulf of Mexico OCS Lease Sale Final Supplemental Environmental Impact Statement 2018 (2018 SEIS) (OCS EIS/EA BOEM 2017-074);
- Biological Opinion Oil and Gas Leasing, Exploration, Development, Production, Decommissioning, and All Related Activities in the Gulf of Mexico Outer Continental Shelf (FWS 2018 BO) (Issued by United States Fish and Wildlife Service [FWS] April 20, 2018); and
- Biological Opinion on the Federally Regulated Oil and Gas Program Activities in the Gulf of Mexico (NMFS 2020 BO) (Issued by National Marine Fisheries Service [NMFS] March 13, 2020).

**Proposed Action:** Anadarko Petroleum Corporation's (Anadarko) Initial Exploration Plan for drilling operations on the OCS of the GOM proposes to explore for hydrocarbons by drilling and completing 36 exploratory wells: Wells A, AA, B, BB, C, CC, D, DD, E, EE, F, FF, G, GG, H, HH, I, II, J, JJ, K, KK, L, LL, M, MM, N, NN, O, OO, P, PP, Q, QQ, R, and RR on Mississippi Canyon Block 80, Lease Number OCS-G35311, Mississippi Canyon Block 81, Lease Number OCS-G35312, Mississippi Canyon 82, Lease Number OCS-G35313, Mississippi Canyon Block 125, Lease Number OCS-G35316, and Mississippi Canyon Block 126, Lease Number OCS-G18194 in the Central Planning Area of the GOM. The proposed activities are located south of Mobile, Alabama approximately 53 miles (85 kilometers) from the nearest shoreline in Plaquemines Parish, Louisiana. The water depths at the proposed well sites range from 3,710 to 4,350 feet (1,131 to 1,326 meters). Anadarko proposes using a dynamically positioned (DP) drillship or DP semisubmersible as the mobile offshore drilling unit (MODU) to drill these wells. Anadarko proposes reduced fuel usage for the DP drillrig and/or DP semisubmersible, and support vessels.

**Resources and Impacts Considered:** The impact analysis for the proposed activity focused on the exploration activities and the resources that may be potentially impacted. The impact producing factors (IPF) include: (1) air emissions; (2) drilling and overboard discharges; (3) seafloor disturbance from well emplacement; (4) vessel traffic and noise; (5) marine trash and debris and other accidental events including vessel strikes, trash, and oil spill.

In this SEA BOEM has considered three alternatives: (1) No Action; (2) Proposed Action as Submitted; and (3) Proposed Action with Conditions of Approval. BOEM has assessed the potential impacts of the proposed action on the following resources:

- 1) air quality;
- 2) water quality;
- 3) deepwater benthic communities;
- 4) marine mammals;
- 5) sea turtles;
- 6) fish resources and essential fish habitat; and
- 7) archaeological resources.

Potential impacts on these resources are summarized below. Direct contact is potentially the most disruptive impact for resources fixed or lying on the sea bottom, and it is weighted most heavily out of all other potentially impacting factors. Pre-activity surveys of the sea bottom required by BOEM may identify

potentially sensitive benthic communities and archaeological resources. At this time no deepwater benthic communities or archaeological resources on the sea bottom are known that could be disturbed by the proposed activity. In the event that either type of resource is encountered, the operator is instructed to avoid impacts to these resources and notify BOEM per the regulations. By operators following the regulations and the regulatory guidance found in the notices to lessees and operators, lease stipulations, and mitigation measures in the NMFS 2020 BO and FWS 2018 BO, potential impacts to air quality, water quality, marine mammals, sea turtles, fish resources and essential fish habitat, and archaeological resources from the proposed activities were determined to be negligible and BOEM will require additional conditions of approval.

Our evaluation in this SEA has selected Alternative 3 and serves as the basis for approving the proposed action. BOEM concludes that no significant impacts are expected to occur to any affected resources by allowing the proposed action to proceed, provided that the specific conditions of approval identified below are met by the operator. Alternative 3 includes the Proposed Action as Submitted with additional mitigation and monitoring measures as required under the Endangered Species Act (ESA) Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO). BOEM concludes that no potentially significant adverse impacts are expected to occur to any affected resources by allowing the proposed action to proceed, provided that the specific mitigation and monitoring measures as part of the ESA Consultation process with NMFS as identified below are met by the operator.

- **COMPLIANCE WITH BIOLOGICAL OPINION TERMS AND CONDITIONS AND REASONABLE AND PRUDENT MEASURES:** This approval is conditioned upon compliance with the Reasonable and Prudent Measures and implementing Terms and Conditions of the Biological Opinion (BO) issued by the National Marine Fisheries Service on March 13, 2020. This includes mitigation, particularly any appendices to Terms and Conditions applicable to the plan, as well as record-keeping and reporting sufficient to allow BOEM and BSEE to comply with reporting and monitoring requirements under the BO; and any additional reporting required by BOEM or BSEE developed as a result of BO implementation. The NMFS BO may be found here:

<https://www.fisheries.noaa.gov/resource/document/biological-opinion-federally-regulated-oil-and-gas-program-activities-gulf-mexico>.

The Appendices and protocols may be found here:

<https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.

- **SUPPORT BASES AND VESSEL TRANSIT ROUTES:** Approval of your plan is conditioned upon your use of the support bases and vessel transit routes as described in your plan. BOEM/BSEE must be notified at least 15 days prior to any vessel route changes that require transit of the Bryde's Whale area, and you must receive prior approval for that transit from BOEM/BSEE.
- **SEISMIC SURVEY OPERATION, MONITORING, AND REPORTING GUIDELINES:** The applicant will follow the guidance provided under Appendix A: Seismic Survey Mitigation and Protected Species Observer Protocols found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on NOAA Fisheries internet website at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.
- **MARINE TRASH AND DEBRIS AWARENESS AND ELIMINATION:** The applicant will follow the guidance provided under Appendix B. Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on NOAA Fisheries internet website at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.
- **VESSEL-STRIKE AVOIDANCE/REPORTING:** The applicant will follow the guidance provided under Appendix C. Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on the NOAA Fisheries

internet site at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.

- **SEA TURTLE RESUSCITATION GUIDELINES:** The applicant will follow the guidance provided under Appendix J. Sea Turtle Handling and Resuscitation Guidelines found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on the NOAA Fisheries internet site at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.
- **MOON POOL MONITORING AND REPORTING:** A moon pool has been identified during review of your plan submittal. If any sea turtle or other marine mammal is detected, you are required to contact NMFS at [nmfs.psoreview@noaa.gov](mailto:nmfs.psoreview@noaa.gov) and BSEE at [protectedspecies@bsee.gov](mailto:protectedspecies@bsee.gov) within 24 hours for additional guidance and incidental report information.

**In addition to the measures proposed by the operator and BSEE/BOEM, the following measures are required by NMFS:**

Application of these measures includes, but is not limited to:

- dive support vessels, service vessels, pipelaying vessels, drillships, floating platforms (e.g., SPAR), mobile offshore drilling units, and other facilities with enclosed moon pools (well in the hull of a vessel, with or without a door).

Moon pool range of activities expected are:

- Deploy or retrieve Remotely Operated Vehicles tethered to vessel
- Deploy or retrieve Autonomous Underwater Vehicle (AUV) tethered to vessel
- Drilling apparatus deployment and retrieval
- Various submarine tools that are attached via tethers
- Use by human divers for entry and exit
- Pipelaying or decommissioning activities

Should your moon pool activities fall outside the scope of those described in the bullets above, then a detailed description of those activities will need to be provided to BOEM and NMFS for review, so that a determination of potential effects to ESA listed species can be made.

- Moon pools with hull doors should attempt to keep doors closed when no activity is occurring within the moon pool, unless the safety of crew or vessel require otherwise. This will prevent ESA-listed species from entering the confined area.
- Use of a moon pool requires regular monitoring while open to the water column and if a vessel is not underway.
- Regular monitoring means 24-hour video monitoring with hourly recurring checks for at least five minutes of the video feed, or, hourly recurring visual checks of the moon pool for at least five minutes by a dedicated crew observer with no other tasks during that visual check.
- If water conditions are such that observers are unable to see within a meter of the surface, operations requiring the lowering or retrieval of equipment through the moon pool must be conducted at a rate that will minimize potential harm to ESA-listed species, if safety permits.

#### **Closure of the hull door**

- Should the moon pool have a hull door that can be closed, then prior to and following closure, the moon pool must be monitored continuously by a dedicated crew observer with no other tasks to ensure that no individual ESA-listed species is trapped within the moon pool. If visibility is not clear to the hull door from above (e.g., turbidity or low light), 30 minutes of monitoring is required prior to hull door closure.

#### **Movement of the vessel (without closed hull door) and equipment deployment/retrieval**

- Prior to movement of the vessel and/or deployment/retrieval of equipment, the moon pool must be monitored continuously for a minimum of 30 minutes, by a dedicated crew observer with no other tasks, to ensure no individual ESA-listed species are present in the moon pool area.

- If an ESA-listed species is observed in the moon pool prior to movement of the vessel, the vessel must not be moved and equipment must not be deployed or retrieved, to the extent practicable, unless the safety of crew or vessel requires otherwise. NMFS personnel must be contacted immediately according to reporting requirements (below) for species type.
  - If the observed animal leaves the moon pool, the operator may commence activities.
  - Should an ESA-listed species be observed in a moon pool prior to activity commencement, recovery of the animal or other actions specific to the scenario may be required to prevent interaction with the animal. Operators must take such action except at the direction of, and after contact with NMFS.
- **SLACK-LINE PRECAUTIONS AND REPORTING REQUIREMENT:**  
**NMFS Requires the following:** If operations require the use of flexible, small diameter (< 2 inch) lines to support operations (with or without divers), operators/contractors must reduce the slack in the lines, to the extent practicable, to prevent accidental entanglement of ESA-listed species. This may include tether lines attached to remotely operated equipment. The following measures are required (noting that diver safety is paramount, and the following measures must be followed only in cases where they do not jeopardize human safety).
    - Operators must utilize tensioning tools and/or other appropriate procedures to reduce unnecessary looseness in the lines and/or potential looping.
    - The lines must remain taut, as long as additional safety risks are not created by this action.
    - A line tender must be present at all times during dive operations and must monitor the line(s) the entire time a diver is in the water.
    - Should the line tender and/or diver become aware of an entanglement of an individual ESA-listed species, the following protocols must be followed as soon as safety permits.

#### **REPORTING REQUIREMENTS:**

Interactions with ESA-listed species must be reported to NMFS and BSEE. Incidents requiring reporting, appropriate reporting contacts, and minimum reporting information are described below. Should any of the following occur at any time, **immediate reporting** of the incident is required (after personnel and/or diver safety is ensured):

- Entanglement or entrapment (i.e., an animal is entangled in a line or cannot or does not leave a moon pool of its own volition) of an ESA-listed species.
- Injury of an ESA-listed species (e.g., the animal appears injured or lethargic).
- Interaction, or contact with equipment by an ESA-listed species.

Contact information for reporting is as follows:

- **Marine mammals:** contact **Southeast Region's Marine Mammal Stranding Hotline at 1-877-433-8299**. If you do not receive a response, go to the following website to contact the relevant stranding networks for marine mammals: <https://www.fisheries.noaa.gov/report>.
- **Sea turtles:** contact **Brian Stacy, Veterinary Medical Officer at 352-283-3370**.
- **Other ESA-listed species** (e.g., giant manta ray, oceanic whitetip shark, or Gulf sturgeon): contact the **ESA Section 7 biologist at 301-427-8413** ([nmfs.psoreview@noaa.gov](mailto:nmfs.psoreview@noaa.gov)).
- **Report all incidents** to [takereport.nmfsser@noaa.gov](mailto:takereport.nmfsser@noaa.gov).
- **Any observation of a leatherback sea turtle** within a moon pool, (regardless of whether it appears injured, or an interaction with equipment or entanglement/entrapment is observed), must be reported immediately to **Brian Stacy, Veterinary Medical Officer at 352-283-3370**.

After the appropriate contacts have been made for guidance/assistance, you may call BSEE at 985-722-7902 for questions or additional guidance on recovery assistance needs (if still required) and continued monitoring requirements.

- Minimum reporting information is described below.
  1. Time, date, water depth, and location (latitude/longitude) of the first discovery of the animal;
  2. Name, type, and call sign of the vessel in which the event occurred;
  3. Equipment being utilized at time of observation;

4. Species identification (if known) or description of the animal involved;
5. Approximate size of animal;
6. Condition of the animal during the event;
7. Photographs or video footage of the animal;
8. NMFS liaison or stranding hotline that was contacted for assistance; and
9. General narrative and timeline describing the events that took place.

**Conclusion:** BOEM has evaluated the potential environmental impacts of the proposed action. Based on SEA No. N-10117 a determination is made that the proposed action would have no significant impact on the marine, coastal, or human environment provided that the avoidance measures required by the specific conditions of approval are met by the operator; therefore, an Environmental Impact Statement will not be required.

---

Chief, Environmental Operations Section  
Office of Environment  
New Orleans Office  
Bureau of Ocean Energy Management

---

October 2, 2020

Date

# TABLE OF CONTENTS

	Page
FINDING OF NO SIGNIFICANT IMPACT (FONSI).....	i
1. OVERVIEW .....	1
1.1. Background.....	2
1.2. Purpose of and Need for the Proposed Action.....	2
1.3. Description of the Proposed Action.....	3
1.4. Impact-Producing Factors.....	3
2. ALTERNATIVES CONSIDERED .....	9
2.1. No Action Alternative.....	9
2.2. Proposed Action as Submitted.....	9
2.3. Proposed Action with Additional Conditions of Approval.....	9
2.4. Summary and Comparison of the Alternatives.....	9
2.5. Alternatives Considered but Not Analyzed.....	13
3. DESCRIPTION OF THE AFFECTED ENVIRONMENT AND ENVIRONMENTAL IMPACTS...	13
3.1. Introduction.....	13
3.2. Air Quality .....	15
3.2.1. Affected Environment.....	15
3.2.2. Impact Analysis.....	16
3.2.2.1. Alternative 1.....	16
3.2.2.2. Alternative 2.....	16
3.2.2.3. Alternative 3.....	17
3.3. Offshore Water Quality.....	17
3.3.1. Affected Environment.....	17
3.3.2. Impact Analysis.....	19
3.3.2.1. Alternative 1.....	19
3.3.2.2. Alternative 2.....	19
3.3.2.3. Alternative 3.....	21
3.4. Deepwater Benthic Communities .....	21
3.4.1. Affected Environment.....	21
3.4.2. Impact Analysis.....	22
3.4.2.1. Alternative 1.....	22
3.4.2.2. Alternative 2.....	22
3.4.2.3. Alternative 3.....	24
3.5. Marine Mammals.....	24
3.5.1. Affected Environment.....	24
3.5.2. Impact Analysis.....	25
3.5.2.1. Alternative 1.....	25
3.5.2.2. Alternative 2.....	26
3.5.2.3. Alternative 3.....	31
3.6. Sea Turtles .....	31
3.6.1. Affected Environment.....	31
3.6.2. Impact Analysis.....	31
3.6.2.1. Alternative 1.....	31
3.6.2.2. Alternative 2.....	32
3.6.2.3. Alternative 3.....	35

3.7.	Fish Resources and Essential Fish Habitat.....	35
3.7.1.	Affected Environment .....	35
3.7.2.	Impact Analysis.....	36
3.7.2.1.	Alternative 1.....	37
3.7.2.2.	Alternative 2.....	37
3.7.2.3.	Alternative 3.....	38
3.8.	Archaeological Resources.....	39
3.8.1.	Affected Environment .....	39
3.8.2.	Impact Analysis.....	40
3.8.2.1.	Alternative 1.....	40
3.8.2.2.	Alternative 2.....	40
3.8.2.3.	Alternative 3.....	42
4.	CONSULTATION AND COORDINATION .....	42
5.	PUBLIC COMMENT .....	43
6.	REFERENCES .....	43
7.	PREPARERS .....	51
8.	APPENDIX.....	52

**SITE-SPECIFIC ENVIRONMENTAL ASSESSMENT (SEA)  
PREPARED FOR  
ANADARKO PETROLEUM CORPORATION  
INITIAL EXPLORATION PLAN:  
N-10117**

## **1. OVERVIEW**

The purpose of this Site-Specific Environmental Assessment (SEA) is to determine whether the proposed activities outlined in the Initial Exploration Plan (EP), N-10117, submitted by Anadarko Petroleum Corporation (Anadarko) on July 10, 2020 will significantly affect the quality of the human environment within the meaning of Section 102(2)(c) of the National Environmental Policy Act (NEPA) and whether an Environmental Impact Statement (EIS) must be prepared. Anadarko’s Initial EP proposes to explore for hydrocarbons by drilling and completing 36 exploratory wells: Wells A, AA, B, BB, C, CC, D, DD, E, EE, F, FF, G, GG, H, HH, I, II, J, JJ, K, KK, L, LL, M, MM, N, NN, O, OO, P, PP, Q, QQ, R, and RR on Mississippi Canyon Block 80, Lease Number OCS-G35311; Mississippi Canyon Block 81, Lease Number OCS-G35312; Mississippi Canyon Block 82, Lease Number OCS-G35313; Mississippi Canyon Block 125, Lease Number OCS-G35316; and Mississippi Canyon Block 126, Lease Number OCS-G18194 in the Central Planning Area (CPA) of the Gulf of Mexico (GOM).

This SEA is tiered from the current NEPA documents that evaluated a broad spectrum of potential impacts resulting from drilling activities across the GOM Outer Continental Shelf (OCS) that include:

- Gulf of Mexico OCS Oil and Gas Lease Sales: 2017-2022 Gulf of Mexico Lease Sales 249, 250, 251, 252, 253, 256, 257, 259, and 261 Final Environmental Impact Statement (Multisale EIS) (USDOJ, BOEM, 2017a);
- Gulf of Mexico OCS Lease Sale Final Supplemental Environmental Impact Statement 2018 (2018 SEIS) (USDOJ, BOEM, 2017b);
- Biological Opinion Oil and Gas Leasing, Exploration, Development, Production, Decommissioning, and All Related Activities in the Gulf of Mexico Outer Continental Shelf (FWS 2018 BO) (Issued by United States Fish and Wildlife Service [FWS] April 20, 2018); and
- Biological Opinion on the Federally Regulated Oil and Gas Program Activities in the Gulf of Mexico (NMFS 2020 BO) (Issued by National Marine Fisheries Service [NMFS] March 13, 2020).

The “tiering” process is provided for in the NEPA implementing regulations (40 CFR § 1502.20 and § 1508.28) and is designed to reduce and simplify the size of subsequent environmental analyses of actions included within the broader program previously examined in NEPA compliance documents by eliminating discussions of impacts that would be repetitive. This allows focus on those site-specific concerns and effects related to the specific action proposed. Document tiering in the Bureau of Ocean Energy Management (BOEM) is subject to additional guidance under the United States Department of the Interior (DOI) regulations at 43 CFR § 46.140 wherein the site-specific analysis must note which conditions and effects addressed in the programmatic document remain valid and which conditions and effects require additional review.

Although the analyses of drilling-related impacts prepared in the Multisale EIS and 2018 SEIS are comprehensive, new information has become available with respect to the following:

- **Emission Impacts on Air Quality** – the EP contains project-specific emissions data not known during the preparation of the programmatic analyses;
- **Discharge Impacts on Offshore Water Quality** – the EP contains project-specific discharge data not known during the preparation of the programmatic analyses;
- **Bottom Impacts on Deepwater Benthic Communities** – the EP contains project-specific information not known during the preparation of the programmatic analyses;

- **Noise/Vessel-Strike Impacts on Marine Mammals** – the environmental baseline since completion of the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO may have experienced slight changes and new information has become available since the preparation of the programmatic analyses;
- **Noise/Vessel-Strike Impacts on Sea Turtles** – the environmental baseline since completion of the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO may have experienced slight changes and new information has become available since the preparation of the programmatic analyses;
- **Discharge Impacts/Disturbances to Fish and Fisheries** – the environmental baseline since completion of the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO may have experienced slight changes and new information has become available since the preparation of the programmatic analyses; and
- **Bottom Impacts on Potential Archaeological Resources** – the EP contains project-specific information not known during the preparation of the programmatic analyses.

Therefore, **Chapter 3** of this SEA focuses on how the new information, including a discussion of the known effects of the *Deepwater Horizon* explosion, spill, and response activities on the analyzed resources, relates to the routine, accidental, and cumulative environmental effects of this proposed action. Where applicable, relevant affected environment discussions and impact analyses from the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO are summarized and utilized for this site-specific analysis, and are incorporated by reference into this SEA. Relevant condition(s) of approval identified in this SEA, Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO have been considered in the evaluation of the proposed action.

## 1.1. BACKGROUND

BOEM and the Bureau of Safety and Environmental Enforcement (BSEE) are mandated to manage and oversee the exploration and development of OCS oil, gas, and mineral resources while ensuring safe operations and the protection of the human, marine, and coastal environments. BOEM and BSEE issue oil and gas leases and regulate exploration, development, production, and decommissioning. Prior to authorizing activities related to these phases, BOEM conducts the appropriate NEPA review. BOEM's Office of Leasing and Plans oversees the submittal of EPs and Development Operations Coordination Documents (DOCD) pursuant to 30 CFR § 550, subpart B.

Lessees and operators submit EPs and DOCDs to provide BOEM with information needed to adequately evaluate the overall potential impacts on OCS resources prior to seeking any individual permit approvals, such as an application for permit to drill (APD). Most of the information in EPs and DOCDs is presented in basic statements, figures, lists, and tables that simply provide the necessary details on the proposed exploration, development, production, and/or transportation operations. One exception is the Environmental Impact Analyses (EIA) required in EPs under 30 CFR § 550.227 and in DOCDs under 30 CFR § 550.261; wherein, the operator provides environmental information and makes impact conclusions regarding their activities.

The scope of the effects on the environment in the GOM from the activities proposed in Anadarko's EP were fully discussed and analyzed in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO and the specific locations, equipment, methodologies, and the duration of the proposed activities will result in impacts similar to those discussed in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO. This SEA was prepared by BOEM to evaluate the activity-specific issues related to the applicant's proposed activities in addition to the new information.

## 1.2. PURPOSE OF AND NEED FOR THE PROPOSED ACTION

Anadarko has submitted a plan to conduct exploration activities on the OCS. The purpose of the proposed action is to drill and complete 36 wells so that Anadarko can utilize the information to evaluate the potential for, and develop plans for, the development and production of hydrocarbon resources on the OCS, which would help satisfy the Nation's need for energy.

The need for this action is established by BOEM's responsibility under the Outer Continental Shelf Lands Act (OCSLA) to make OCS lands available for expeditious and orderly development, subject to

environmental safeguards, in a manner that is consistent with the maintenance of competition and other national needs. Section 11 of OCSLA (43 U.S.C. § 1340) requires oil and gas lessees seeking to conduct exploration activities to first obtain approval from the Secretary who has delegated the authority to grant such approval to BOEM.

In response to the proposed action in Anadarko's plan, BOEM is required by OCSLA to approve, approve with modifications, or deny the plan within 30 days (see 43 U.S.C. § 1340(c)(1)). The criteria that BOEM will apply in reaching a decision to approve, approve with modifications, or deny the plan within 30 days and the scope of its discretion are provided by Section 11 of OCSLA and detailed in the implementing regulations (30 CFR § 550, subpart B). Authorizing the proposed action, as outlined in the Initial EP N-10117, allows Anadarko to pursue its rights under the lease and to conduct exploration drilling activities.

### 1.3. DESCRIPTION OF THE PROPOSED ACTION

Anadarko's Initial EP for drilling operations on the OCS of the GOM proposes to explore for hydrocarbons by drilling and completing 36 exploratory wells: Wells A, AA, B, BB, C, CC, D, DD, E, EE, F, FF, G, GG, H, HH, I, II, J, JJ, K, KK, L, LL, M, MM, N, NN, O, OO, P, PP, Q, QQ, R, and RR on Mississippi Canyon Block 80, Lease Number OCS-G35311; Mississippi Canyon Block 81, Lease Number OCS-G35312; Mississippi Canyon Block 82, Lease Number OCS-G35313; Mississippi Canyon Block 125, Lease Number OCS-G35316; and Mississippi Canyon Block 126, Lease Number OCS-G18194 in the Central Planning Area of the GOM. The proposed activities are located south of Mobile, Alabama approximately 53 miles (mi) (85 kilometers [km]) from the nearest shoreline in Plaquemines Parish, Louisiana. The water depths at the proposed well sites range from 3,710 to 4,350 feet (ft) (1,131 to 1,326 meters [m]). Anadarko proposes using a dynamically positioned (DP) drillship or DP semisubmersible as the mobile offshore drilling unit (MODU) to drill these wells. Anadarko proposes reduced fuel usage for the drillrig(s) and support vessels. The projected duration of the proposed drilling and completing for one well is 85 days; with proposed drilling planned from January 2021 to September 2030.

Supply, support, and crewboat facilities to support the proposed action are to be located in existing facilities in Port Fourchon, Louisiana, approximately 121 mi (195 km) northwest of the project location. Port Fourchon will be used as the primary debarkation point for equipment, supplies, and crews supporting the proposed action. Air operations to support the proposed action are to be located from existing facilities in Houma, Louisiana, approximately 157 mi (253 km). Anadarko does not expect any shore-based construction or expansion in association with this proposed action. The types of support vessels and their potential travel frequency during exploratory drilling are included Anadarko's plan (Anadarko, 2020).

### 1.4. IMPACT-PRODUCING FACTORS

An impact-producing factor (IPF) is any activity or process resulting from an approved operation that causes impacts to the environment, such as an emission, effluent, or physical disturbance. The IPFs from the routine activities proposed by the operator in this plan include: (1) waste and discharges from vessel operations and exploration activities; (2) air emissions from equipment and vessels; (3) noise from vessel and helicopter transportation and drilling activities; and (4) bottom disturbances from well emplacement activities. The routine IPFs are expected to occur during the operations conducted under the proposed action and are addressed in each of the site-specific analyses in **Chapter 3** under "Routine Activities."

The analyses in **Chapter 3** also consider IPFs that might result from an accidental event. The primary IPFs from potential accidents related to OCS drilling activities include: (1) vessel collisions with marine mammals and sea turtles; (2) oil spills and blowouts; (3) bottom disturbances from lost/jettisoned debris; and (4) helicopter collisions with coastal and/or marine birds. Unlike the IPFs associated with routine activities, the IPFs from accidental events are not expected because of the low probabilities of such events from occurring, existing/recently implemented safety measures and condition(s) of approval, and an increased level of operator awareness observed since the *Deepwater Horizon* spill. The accidental IPFs are detailed and addressed in each of the site-specific analyses under "Accidental Events."

The Multisale EIS and 2018 SEIS considered the routine and accidental IPFs described above; however, additional information related to the oil spill/blowout IPF has been collected since the *Deepwater Horizon* spill that was not available during the preparation of the programmatic analyses. [Appendix A; Accidental Oil-Spill Discussion \(http://www.boem.gov/Appendix-A-Deepwater-SEA-Oil-Spill-Discussion/\)](http://www.boem.gov/Appendix-A-Deepwater-SEA-Oil-Spill-Discussion/),

introduces the new data and describes the circumstances that might result if an accidental spill were to occur. Additionally, the analyses of the “Accidental Events” incorporate information from Appendix B of the 2012-2017 WPA/CPA Multisale EIS (USDOJ, BOEM, 2012), “Catastrophic Spill Event Analysis,” to address the potential impacts to the environment in the unlikely event that a catastrophic spill similar to the *Deepwater Horizon* spill was to occur. This analysis was later updated and published as a “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c) and incorporated by reference.

**Accidental Spill Concerns**

Since spills are unplanned, unforeseeable events, BOEM is required to rely on past experiences to predict many factors regarding oil-spill risks. Based on experience and the operations proposed in Anadarko’s plan, the potential sources of hydrocarbon spills from the proposed activity would include the following:

- A storage tank accident on the MODU;
- A transfer operation mishap between the supply vessel and the MODU;
- A leak resulting from damage to the fuel tanks on one of the supply or crew boats; and/or
- A blowout of a proposed well.

***Potential Spills from Vessels/Transfer Operations***

As indicated above, offshore spills from Anadarko’s proposed action are possible if an accident were to damage a storage tank onboard the drilling rig, the crewboat, offshore support vessel, or the fuel supply vessel. Historically, accidents of this nature have resulted from unintentional vessel collisions and transfer incidents during the offloading of diesel fuel to the drilling rig. Anadarko plans to use a DP drillship or DP semisubmersible as the MODUs with a subsea blowout preventer (BOP) to conduct the proposed activities. There are several tanks onboard the MODU that store fuel and lubricants necessary for the rig’s operation. A worst-case discharge scenario from a rupture and spill from the vessels are:

	<u>Largest Main Tank Capacity</u>	<u>Total Diesel Oil Capacity</u>
DP Semisubmersible	4,541 bbl	16,689 bbl
DP Drillship	12,458 bbl	62,874 bbl

Additionally, the support vessels proposed to aid the drilling operations have an estimated fuel tank capacity of 450,698 gallons (gal) (10,731 bbl), the crew boat has an estimated fuel tank capacity of 70,000 gal (1,667 bbl), and the supply and diesel oil supply vessels have an estimated fuel tank capacity of 336,227 gal (8,005 bbl). The helicopter proposed to support activities has an estimated fuel tank capacity of 735 gal (18 bbl) (Anadarko, 2020).

***Potential Spills from a Loss of Well Control/Blowout***

BOEM and BSEE require that all losses of well control (blowouts) be reported. The current definition for “loss of well control” used by BOEM and BSEE is:

- Uncontrolled flow of formation or other fluids. The flow may be to an exposed formation (an underground blowout) or at the surface (a surface blowout);
- flow through a diverter; or
- uncontrolled flow resulting from a failure of surface equipment or procedures.

Losses of well control (also known as blowouts) can occur during exploratory drilling, development drilling, completion, production, or workover operations. A blowout can occur when improperly balanced well pressures result in the sudden, uncontrolled releases of fluids from a wellbore or wellhead (PCCI, 1999; Neal Adams Firefighters, Inc., 1991). Since 1971, most OCS blowouts have resulted in the release of gas; blowouts resulting in the release of oil have been rare. The most recent blowout was related to the *Deepwater Horizon* spill, which resulted in the release of both gas and oil. In the event of a blowout, an operator’s first course of action is to activate the BOP to close the well. The BOP may be located on the

surface of the drilling rig or on the seafloor. There are built-in redundancies in the BOP system to allow activation of selected components with the intent to seal off the well bore. If a subsea BOP cannot be operated from the drill rig, it can be operated at the seafloor using remotely operated vehicles (ROV).

If the blowout occurs during drilling, pieces of the rock formation below the drill bit may fail and collapse into the wellbore because of the pressure drop. Formation fragments subsequently clog or “bridge” the drill bit or pipe, reducing or stopping flow (PCCI, 1999). Completed wells, or those in production, present more severe consequences in the event of a blowout due to the hole being fully cased down to the producing formation that lowers the probability of bridging (PCCI, 1999).

If the BOP fails and the well does not bridge, there are other options available to control the blowout that include capping/shut-in, capping/diverting, surface stinger, vertical intervention, offset kill, and drilling relief wells (Neal Adams Firefighters, Inc., 1991). Of these methods, a relief well is the most important remedy and may be required immediately (even if it is not the first choice), since it is typically considered the ultimate solution for well control. A relief well must be drilled from a nearby platform or drillship. It is estimated that drilling a relief well in deep water can take anywhere from 30 to 120 days or more. The actual amount of time required to drill the relief well will depend upon the complexity of the intervention, the location of a suitable rig, the type of operation that must be terminated in order to release the rig (e.g., may need to secure the well before releasing the rig), and any problems mobilizing personnel and equipment to the location.

### ***Catastrophic Spill Event Analysis***

After the *Deepwater Horizon* explosion, oil spill, and response, BOEM prepared a “Catastrophic Spill Event Analysis,” a region-wide evaluation that identifies the most likely and most significant impacts from a high-volume blowout and oil spill that continues for an extended period of time, which was included as Appendix B in the 2012-2017 WPA/CPA Multisale EIS (USDOJ, BOEM, 2012). Since then, this analysis has been reviewed and updated. This analysis, which is based on credible scientific evidence, identifies the most likely and most significant impacts from a high-volume blowout and oil spill that continues for an extended period of time and has been published as an independent white paper, “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c). The scenario and impacts discussed in that white paper are comparable to that of a spill similar to the *Deepwater Horizon* spill and are not associated with IPFs anticipated to result from routine activities or even more reasonably feasible, accidental events that could occur during the proposed action. The conclusions made in that white paper are addressed in the SEA’s impact analyses (**Chapters 3.2 to 3.8**) and incorporated by reference.

### ***Site-Specific Estimate of Spill Risk***

Anadarko’s plan describes measures for blowout prevention, likelihood for surface intervention to stop a blowout, and early intervention in the event of a blowout. Anadarko has developed standards for well control, personnel safety, and an emergency response plan and these methods are stated in detail in the plan submitted by Anadarko (Anadarko, 2020). As per the information provided in Anadarko’s plan, the MODU Anadarko plans to use will deploy a subsea BOP while drilling the well (Anadarko, 2020). An estimate of spill risk from Anadarko’s proposed activities was calculated using the drilling spill rate for the entire OCS and the estimated number of wells to be drilled. The resulting value, 0.00252 or 0.252 percent, is used to address the risk of a spill >1,000 bbl occurring during the proposed action. When examining only wells in deep water (in water depths >500 ft; [152 m]), past data suggest the chance of a major spill from a deepwater well under current regulations and practices is 1 in 4,957 (USDOJ, BOEM, 2012).

Though not proposed or expected, Anadarko has estimated that a worst-case discharge (WCD) scenario from a blowout of one of the well under this proposed action could be 371,735 BOPD of 33.1° American Petroleum Institute (API) gravity crude. In accordance with enhanced agency oversight, BOEM verified the operator’s calculations used to determine the WCD volume.

The WCD scenario does not include any bridging. Mechanical collapse of the reservoirs in the openhole section of the wellbore was not considered in determining WCD. During a WCD event, the openhole portion of the well will be exposed to a substantial underbalance condition. Due to the unconsolidated nature of the formations contributing flow and the relatively weak remaining exposed sediments, a significant quantity sand and heaving shale will enter the flowstream. The presence of sediments in the flowstream are excluded from Anadarko’s discharge calculations and assumes no bridging will occur; however, bridging is likely to occur. Anadarko has developed standards for well control,

personnel safety, and emergency response plan. These methods are stated in detail in the plan (Anadarko, 2020).

In the event a relief well is required due to blowout, Anadarko indicates in their plan that through Mutual Aid agreements rigs that are capable of operating at water and reservoir depths similar to the proposed activities are available (Anadarko, 2020). For this project, Anadarko estimates that it will take approximately 7 to 21 days to secure a well, demobilize the drill rig, and transit to the relief well site; 60 to 70 days to drill the relief well down to the target zone, for a total of 67 to 91 days to drill a relief well. Also, there are no existing facilities/platforms nearby from where the relief well can be drilled. Additional details related to the proposed action can be found in Anadarko's proposed exploration plan (Anadarko, 2020).

### ***Spill Response Requirements***

Agency regulations require that all owners and operators of oil handling, storage, or transportation facilities located seaward of the coastline submit an Oil Spill Response Plan (OSRP) before they can use a facility. BSEE has issued notices to lessees and operators (NTL) No. 2012-N06 (*Guidance to Owners and Operators of Offshore Facilities Seaward of the Coast Line Concerning Regional Oil Spill Response Plans*), which informs operators of OSRP requirements and requires that they have adequate resources available to protect the environment from spills from their facilities. The Environmental Protection and Response Plan within the OSRP outlines the availability of spill containment and cleanup equipment and trained personnel necessary to ensure that a full response can be deployed during an oil spill emergency. All the proposed activities and facilities in this plan will be covered by the Regional OSRP O-16 filed by Anadarko Petroleum Corporation and its subsidiary Anadarko US Offshore LLC (Operator Numbers 00689 and 02219) in accordance with 30 CFR 550 and 30 CFR 254 and deemed in compliance by BSEE with the last biannual update found to be in compliance in September 2019. Anadarko also certifies it has the capability to respond, to the maximum extent practicable, to worst-case discharge, or a substantial threat of such a discharge, resulting from the activities proposed in their Initial EP (Anadarko, 2020).

### **Spill Response, Oil Spill Response Plan, BSEE Spill Response Program**

A discussion about spill response, the Oil Spill Response Plan, and the BSEE Spill Response Program is already included in Appendix A: <http://www.boem.gov/Appendix-A-Deepwater-SEA-Oil-Spill-Discussion/>.

### ***Subsurface Response***

Most oil-spill response strategies and equipment are based upon the simple principle that oil floats. However, as evident during the *Deepwater Horizon* spill, this is not always true. Sometimes oil suspends within the water column or sinks to the seafloor and sometimes it does all three: floats, suspends, and sinks. Oil suspended in the water column and moving with the currents is difficult to track using standard visual survey methods. Trajectory models traditionally used to predict floating oil movement and fate are not applicable to submerged oil - oil that is suspended in the water column and/or that sinks. There are no proven methods for the containment of submerged oil, and methods for recovery of submerged oil have limited effectiveness (Coastal Response Research Center, 2007).

Efforts to contain and/or recover suspended oil have focused on different types of nets, either the ad hoc use of fishing nets or specially designed trawl nets. There has been research conducted on the design of trawl nets for recovery of emulsified fuels. However, the overall effectiveness for large spills is expected to be very low. Suspended oil can occur as liquid droplets or semisolid masses in sizes ranging from millimeters to meters in diameter. At spills where oil has been suspended in the water column, responders have devised low technology methods for tracking the presence and spread of oil over space and time. For suspended oil, these methods include stationary systems such as snare sentinels, which can consist of any combination of the following: a single length of snare on a rope attached to a float and an anchor; one or more crab traps on the bottom that are stuffed with snare; and minnow or other type of traps that are stuffed with snare and deployed at various water depths. The configuration would depend upon the water depth where the oil is located within the water column. Currently, it is not possible to determine the particle size, number of particles, or percent oil cover in the water column based upon the visual observations of oil on these systems (Coastal Response Research Center, 2007).

Spills involving submerged oil trigger the need for real-time data on current profiles (surface to bottom), wave energy, suspended sediment concentrations, detailed bathymetry, seafloor sediment characteristics,

and sediment transport patterns and rates. These data are needed to validate or calibrate models (both computer and conceptual), direct sampling efforts, and predict the behavior and fate of the submerged oil. This information might be obtained through the use of acoustic Doppler current profilers, dye tracer studies, rapid seafloor mapping systems, and underwater camera or video systems that could record episodic events (Coastal Response Research Center, 2007). During the *Deepwater Horizon* spill, fluorometers were used successfully to detect the presence of oil.

### ***Surface Response***

Prior to the DeepSpill sea trials, there was some doubt about whether oil released subsea in deep water would reach the sea surface. The surface slick formed after the DeepSpill crude oil releases contained patches of water-in-oil emulsion with film thickness more than adequate for containment with oil booms and also sufficient thickness for efficient treatment with chemical dispersant, similar to what actually happened during the *Deepwater Horizon* spill. However, the DeepSpill sea trials indicated that the potential lifetime of the crude oil slick would be short, which resulted in the report suggesting that the slick could be left to disperse naturally without attempting any mechanical cleanup (Johansen et al., 2001). The fact that the experiment did not involve the quantity of crude that was lost per day and on an ongoing basis for approximately 87 days as occurred during the *Deepwater Horizon* spill may account for the observed differences in slick behavior between the experiments and the GOM spill. As occurred during the Norwegian Sea trials, there was no hydrate formation at the damaged riser during the uncontrolled flow during the *Deepwater Horizon* spill.

The *Deepwater Horizon* spill incident indicated that, although released at a water depth of 5,000 ft (1,524 m), once the oil surfaced, a variety of response methods were effective on the oil that surfaced near the source. The options for oil combat in deep water are the same as those used for shallower waters (mechanical recovery, dispersion, in-situ burning). Response to the oil as it emulsified and moved farther from the source proved more difficult. The emulsified oil had to be chased down by the responders, making it more difficult for the skimmers to stay in skimmable oil. The emulsified oil was also less likely to be effectively burned or dispersed.

A variety of standard cleanup protocols were used for removing *Deepwater Horizon* oil from beaches, shorelines, and offshore water (**Table 1**). After the *Deepwater Horizon* spill, BSEE (then BOEMRE) issued NTL No. 2010-N10 (*Statement of Compliance with Applicable Regulations and Evaluation of Information Demonstrating Adequate Spill Response and Well Containment Resources*) that became effective on November 8, 2010. This NTL applies only to operators conducting operations using subsea blowout BOPs or surface BOPs on floating facilities. The NTL also informs lessees that BSEE will be evaluating whether each operator has submitted adequate information demonstrating that it has access to and can deploy surface and subsea containment resources that would be adequate to promptly respond to a blowout or other loss of well control. Although the NTL does not require that operators submit revised OSRPs that include this containment information at this time, operators were notified of BSEE's intention to evaluate the adequacy of each operator to comply in their current OSRP.

### ***Source Control and Containment***

The type of information that BSEE reviews pursuant to this NTL includes, but is not limited to:

- Subsea containment and capture equipment, including containment domes and capping stacks;
- Subsea utility equipment, including hydraulic power, hydrate control, and dispersant injection equipment;
- Riser systems;
- Remotely operated vehicles;
- Capture vessels;
- Support vessels; and
- Storage facilities.

Table 1  
Primary Cleanup Options Used during the *Deepwater Horizon* Spill Response.

Type	Fresh Oil	Sheens	Mousse	Tar Balls	Burn Residue
On-Water Response	Disperse, skim, burn	Light sheens very difficult to recover, heavier sheens picked up with sorbent boom or sorbent pads	Skim	Snare boom	Manual removal
On-Land Response	Sorbent pads, manual recovery, flushing with water, possible use of chemical shoreline cleaning agents	Light sheens very difficult to recover, heavier sheens picked up with sorbent boom or sorbent pads	Sorbent pads, manual recovery	Snare boom, manual removal, beach cleaning machinery	Manual removal

Source: USDOC, NOAA, 2010.

To address the improved containment systems expectations to rapidly contain a spill as a result of a loss of well control from a subsea well (addressed in NTL No. 2010-N10), several oil and gas industry majors initiated the development of a rapid response system. This system is designed to fully contain oil flow in the event of a potential future underwater blowout and to address a variety of scenarios. The system would consist of specially designed equipment constructed, tested, and available for rapid response. It is envisioned that this system could be fully operational within days to weeks after a spill event occurs. The system is designed to operate in up to 10,000 ft (3,048 m) water depth and will add containment capability of 100,000 BOPD (4.2 million gallons per day). The companies that originated this system have formed a non-profit organization, the Marine Well Containment Company (MWCC), to operate and maintain the system. MWCC will provide fully trained crews to operate the system, will ensure the equipment is operational and ready for rapid response and will conduct research on new containment technologies. This system will connect by risers to vessels that are designed to safely capture, store and offload the oil. This improves safety and environmental protection by fully securing the well via capping and shut-in or by containing the oil flow until the well is under control. It also enhances safe operations by reducing congestion (i.e., fewer vessels, risers/flowlines). Until this equipment is available, MWCC has built a subsea containment equipment system that is engineered to be used in water depths up to 8,000 ft (2,438 m) and has the capacity to contain 60,000 BOPD. This initial response system includes a capping stack with the ability to shut in oil flow or to flow the oil via flexible pipes and risers to surface vessels.

Another option for source control and containment is through the use of the equipment stockpiled by Helix Energy Solutions Group, Inc. (Helix). The Helix initiative involves more than 20 smaller energy companies, and supplements the MWCC response effort. Helix has stockpiled the equipment that it found useful in the *Deepwater Horizon* response and is offering it to oil and gas producers for immediate use. The Helix system centers on three ships: the *Helix Producer I*; the *Q4000*; and the *Express* deepwater construction vessel. These vessels played a role in the *Deepwater Horizon* response and continue to work in the Gulf. Together, the Helix ships and related equipment can handle up to 55,000 BOPD, 70,000 bbl of liquid natural gas, and 95 million cubic ft of natural gas at depths up to 8,000 ft (2,438 m). The primary difference between the MWCC system and the Helix system is that nothing needs to be built for the Helix system; it has been field tested and is currently available for deployment. Another group, Wild Well Control, is also providing some subsea containment capability and debris removal to offshore operators.

BOEM and BSEE will not allow an operator to begin drilling operations until adequate subsea containment and collection equipment as well as subsea dispersant capability is determined by the bureau

to be available to the operator and sufficient for use in response to a potential incident from the proposed well(s). However, it would be impossible to predict with any degree of certainty the percentage of oil that could be contained subsea in the event of a spill or when or if complete containment would even be possible. There are some situations where this equipment might not be able to be used to control the well, for example, if the drilling structure were to fall directly on top of the well as debris during a loss of well control event. If a loss of well control event occurred in the future, it is possible that it could be contained in a best-case scenario within weeks with the utilization of the rapid subsea containment packages, thereby greatly limiting the amount of oil potentially lost to the environment.

## Summary

In the event of a spill, particularly a blowout, there is no single method of containing and removing it that would be 100 percent effective. Removal and containment efforts to respond to an ongoing spill would likely require multiple technologies, including mechanical cleanup, burning of the slick, and chemical dispersants. Even with the deployment of all of these technologies, it is likely that, with the operating limitations of today's spill response technology, not all of the oil could be contained and removed offshore. It is likely that larger spills in deep waters and under the right conditions would require the simultaneous use of all available cleanup methods (mechanical cleanup, dispersant application, and in-situ burning). That being said, when one considers the historical/statistical data, the recent subsea containment improvements, BOEM's and BSEE's enhanced oversight, and industry's heightened safety awareness since the *Deepwater Horizon* spill, it is reasonable to conclude that an accidental spill event is not likely to occur.

## 2. ALTERNATIVES CONSIDERED

### 2.1. NO ACTION ALTERNATIVE

**Alternative 1** - If selected, the operator would not undertake the proposed activities. If the proposed activities are not undertaken, all environmental impacts, including additional routine, accidental, or cumulative impacts to the environmental and cultural resources described in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO would not occur. Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas exploration and development activities as described in the Multisale EIS and 2018 SEIS and routine, accidental, and cumulative impacts would continue to occur.

### 2.2. PROPOSED ACTION AS SUBMITTED

**Alternative 2** - If selected, the operator would undertake the proposed activities as requested in their plan. This alternative assumes that the operator will conduct their operations in accordance with their lease stipulations (in addition to any amendments), the OCSLA, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), conditions of approval as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and with FWS that concluded on April 20, 2018 (FWS 2018 BO). However, no additional, site-specific mitigation or monitoring measures would be required by BOEM or NMFS.

### 2.3. PROPOSED ACTION WITH ADDITIONAL CONDITIONS OF APPROVAL

**Alternative 3** - This is BOEM's *Preferred Alternative*. If selected, the operator would undertake the proposed activities as requested in the plan and conditioned by stipulations, regulations, and guidance (similar to Alternative 2); however, BOEM and NMFS would require the operator to undertake additional mitigation and monitoring measures as identified in **Chapter 2.4** below and described in the effects analyses in order to fully address the potential site-specific impacts of the proposed activities.

### 2.4. SUMMARY AND COMPARISON OF THE ALTERNATIVES

If selected, Alternative 1, the No Action Alternative, would result in the operator not exercising its rights under the lease and conducting their proposed activities. Alternative 1 would not result in any impacts

to the environmental resources analyzed in **Chapter 3**; however, the lessee would not develop the oil and gas resources of its lease for the benefit of the U.S. economy. Alternative 2 would result in the lessee achieving its objectives; however, BOEM and NMFS have determined that additional mitigation and monitoring measures (described below) are needed to minimize or reduce the risk of possible environmental impacts. Alternative 2 does not include the additional mitigation and monitoring measures from NMFS.

Alternative 3 is the Preferred Alternative because it allows the lessee to achieve its exploration objectives and also provides for additional mitigation and monitoring requirements to minimize or reduce the risk to potential environmental impacts.

### **Condition(s) of Approval Required under the Preferred Alternative**

The need for, and utility of, the following conditions of approval discussed in the relevant impact analysis section of this SEA. Alternative 3 is the Proposed Action with Additional Conditions of Approval as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and NMFS concurrence received on September 28, 2020. To ensure adequate environmental protection, the following conditions of approval are applied:

- **COMPLIANCE WITH BIOLOGICAL OPINION TERMS AND CONDITIONS AND REASONABLE AND PRUDENT MEASURES:** This approval is conditioned upon compliance with the Reasonable and Prudent Measures and implementing Terms and Conditions of the Biological Opinion (BO) issued by the National Marine Fisheries Service on March 13, 2020. This includes mitigation, particularly any appendices to Terms and Conditions applicable to the plan, as well as record-keeping and reporting sufficient to allow BOEM and BSEE to comply with reporting and monitoring requirements under the BO; and any additional reporting required by BOEM or BSEE developed as a result of BO implementation. The NMFS BO may be found here:

<https://www.fisheries.noaa.gov/resource/document/biological-opinion-federally-regulated-oil-and-gas-program-activities-gulf-mexico>.

The Appendices and protocols may be found here:

<https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.

- **SUPPORT BASES AND VESSEL TRANSIT ROUTES:** Approval of your plan is conditioned upon your use of the support bases and vessel transit routes as described in your plan. BOEM/BSEE must be notified at least 15 days prior to any vessel route changes that require transit of the Bryde's Whale area, and you must receive prior approval for that transit from BOEM/BSEE.
- **SEISMIC SURVEY OPERATION, MONITORING, AND REPORTING GUIDELINES:** The applicant will follow the guidance provided under Appendix A: Seismic Survey Mitigation and Protected Species Observer Protocols found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on NOAA Fisheries internet website at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.
- **MARINE TRASH AND DEBRIS AWARENESS AND ELIMINATION:** The applicant will follow the guidance provided under Appendix B. Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on NOAA Fisheries internet website at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.
- **VESSEL-STRIKE AVOIDANCE/REPORTING:** The applicant will follow the guidance provided under Appendix C. Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on the NOAA Fisheries internet site at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.

- **SEA TURTLE RESUSCITATION GUIDELINES:** The applicant will follow the guidance provided under Appendix J. Sea Turtle Handling and Resuscitation Guidelines found in the Biological Opinion issued by the National Marine Fisheries Service on March 13, 2020. The guidance can be accessed on the NOAA Fisheries internet site at <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>.
- **MOON POOL MONITORING AND REPORTING:** A moon pool has been identified during review of your plan submittal. If any sea turtle or other marine mammal is detected, you are required to contact NMFS at [nmfs.psoreview@noaa.gov](mailto:nmfs.psoreview@noaa.gov) and BSEE at [protectedspecies@bsee.gov](mailto:protectedspecies@bsee.gov) within 24 hours for additional guidance and incidental report information.

**In addition to the measures proposed by the operator and BSEE/BOEM, the following measures are required by NMFS:**

Application of these measures includes, but is not limited to:

- dive support vessels, service vessels, pipelaying vessels, drillships, floating platforms (e.g., SPAR), mobile offshore drilling units, and other facilities with enclosed moon pools (well in the hull of a vessel, with or without a door).

Moon pool range of activities expected are:

- Deploy or retrieve Remotely Operated Vehicles tethered to vessel
- Deploy or retrieve Autonomous Underwater Vehicle (AUV) tethered to vessel
- Drilling apparatus deployment and retrieval
- Various submarine tools that are attached via tethers
- Use by human divers for entry and exit
- Pipelaying or decommissioning activities

Should your moon pool activities fall outside the scope of those described in the bullets above, then a detailed description of those activities will need to be provided to BOEM and NMFS for review, so that a determination of potential effects to ESA listed species can be made.

- Moon pools with hull doors should attempt to keep doors closed when no activity is occurring within the moon pool, unless the safety of crew or vessel require otherwise. This will prevent ESA-listed species from entering the confined area.
- Use of a moon pool requires regular monitoring while open to the water column and if a vessel is not underway.
- Regular monitoring means 24-hour video monitoring with hourly recurring checks for at least five minutes of the video feed, or, hourly recurring visual checks of the moon pool for at least five minutes by a dedicated crew observer with no other tasks during that visual check.
- If water conditions are such that observers are unable to see within a meter of the surface, operations requiring the lowering or retrieval of equipment through the moon pool must be conducted at a rate that will minimize potential harm to ESA-listed species, if safety permits.

#### **Closure of the hull door**

- Should the moon pool have a hull door that can be closed, then prior to and following closure the moon pool must be monitored continuously by a dedicated crew observer with no other tasks to ensure that no individual ESA-listed species is trapped within the moon pool. If visibility is not clear to the hull door from above (e.g., turbidity or low light), 30 minutes of monitoring is required prior to hull door closure.

#### **Movement of the vessel (without closed hull door) and equipment deployment/retrieval**

- Prior to movement of the vessel and/or deployment/retrieval of equipment, the moon pool must be monitored continuously for a minimum of 30 minutes, by a dedicated crew observer with no other tasks, to ensure no individual ESA-listed species are present in the moon pool area.
- If an ESA-listed species is observed in the moon pool prior to movement of the vessel, the vessel must not be moved and equipment must not be deployed or retrieved, to the extent

- practicable, unless the safety of crew or vessel requires otherwise. NMFS personnel must be contacted immediately according to reporting requirements (below) for species type.
- If the observed animal leaves the moon pool, the operator may commence activities.
  - Should an ESA-listed species be observed in a moon pool prior to activity commencement, recovery of the animal or other actions specific to the scenario may be required to prevent interaction with the animal. Operators must take such action except at the direction of, and after contact with NMFS.

- **SLACK-LINE PRECAUTIONS AND REPORTING REQUIREMENT:**

**NMFS Requires the following:** If operations require the use of flexible, small diameter (< 2 inch) lines to support operations (with or without divers), operators/contractors must reduce the slack in the lines, to the extent practicable, to prevent accidental entanglement of ESA-listed species. This may include tether lines attached to remotely operated equipment. The following measures are required (noting that diver safety is paramount, and the following measures must be followed only in cases where they do not jeopardize human safety).

- Operators must utilize tensioning tools and/or other appropriate procedures to reduce unnecessary looseness in the lines and/or potential looping.
- The lines must remain taut, as long as additional safety risks are not created by this action.
- A line tender must be present at all times during dive operations and must monitor the line(s) the entire time a diver is in the water.
- Should the line tender and/or diver become aware of an entanglement of an individual ESA-listed species, the following protocols must be followed as soon as safety permits.

**REPORTING REQUIREMENTS:**

Interactions with ESA-listed species must be reported to NMFS and BSEE. Incidents requiring reporting, appropriate reporting contacts, and minimum reporting information are described below. Should any of the following occur at any time, **immediate reporting** of the incident is required (after personnel and/or diver safety is ensured):

- Entanglement or entrapment (i.e., an animal is entangled in a line or cannot or does not leave a moon pool of its own volition) of an ESA-listed species.
- Injury of an ESA-listed species (e.g., the animal appears injured or lethargic).
- Interaction, or contact with equipment by an ESA-listed species.

Contact information for reporting is as follows:

- **Marine mammals:** contact **Southeast Region's Marine Mammal Stranding Hotline at 1-877-433-8299**. If you do not receive a response, go to the following website to contact the relevant stranding networks for marine mammals: <https://www.fisheries.noaa.gov/report>.
- **Sea turtles:** contact **Brian Stacy, Veterinary Medical Officer at 352-283-3370**.
- **Other ESA-listed species** (e.g., giant manta ray, oceanic whitetip shark, or Gulf sturgeon): contact the **ESA Section 7 biologist at 301-427-8413** ([nmfs.psoreview@noaa.gov](mailto:nmfs.psoreview@noaa.gov)).
- **Report all incidents to** [takereport.nmfs@noaa.gov](mailto:takereport.nmfs@noaa.gov).
- **Any observation of a leatherback sea turtle** within a moon pool, (regardless of whether it appears injured, or an interaction with equipment or entanglement/entrapment is observed), must be reported immediately to **Brian Stacy, Veterinary Medical Officer at 352-283-3370**.

After the appropriate contacts have been made for guidance/assistance, you may call BSEE at 985-722-7902 for questions or additional guidance on recovery assistance needs (if still required) and continued monitoring requirements.

- Minimum reporting information is described below.
  1. Time, date, water depth, and location (latitude/longitude) of the first discovery of the animal;
  2. Name, type, and call sign of the vessel in which the event occurred;
  3. Equipment being utilized at time of observation;
  4. Species identification (if known) or description of the animal involved;

5. Approximate size of animal;
6. Condition of the animal during the event;
7. Photographs or video footage of the animal;
8. NMFS liaison or stranding hotline that was contacted for assistance; and
9. General narrative and timeline describing the events that took place.

## 2.5. ALTERNATIVES CONSIDERED BUT NOT ANALYZED

Several other alternatives were considered and reviewed during the preparation of this SEA and coordination of the resource reviews. Ultimately, a viable alternative is required to be a logical option for carrying out the proposed action, ensure that the purpose and need can be met, and be feasible under the regulatory directives of the OCSLA and all other applicable guidance. The table below lists the alternatives that were considered but dismissed and not analyzed further along with the rationale for the decision:

Alternatives Considered but Not Analyzed.

Dismissed Alternative	Alternative Detail	Reason Not Analyzed
Daytime Drilling Only.	The alternative would restrict all drilling operations to the hours between legal sunrise and sunset to take advantage of the increased lighting in an effort to improve safety.	This alternative does not consider that adequate lighting is available on vessels and MODUs, existing safety protocols, and that the premature stopping of some drilling/well operations prior to critical junctures could lead to highly problematic and unsafe situations.
Drilling from an Anchored MODU Only.	The alternative would only allow drilling activities from an anchored MODU to reduce air quality impacts from the increased emissions released from DP MODUs.	This alternative does not consider the limited availability of conventionally moored MODUs in the GOM or the negligible air quality concerns for temporary operations taking place a great distance from shore.
Incorporation of “Seasonal” Drilling Windows.	The alternative would be based upon observed “seasonal” migrations or behavioral patterns exhibited by marine protected species (MPS) and would restrict the proposed drilling operations for several weeks/months each year.	This alternative would have to rely upon incomplete seasonal data as most migratory MPS are not found in the GOM and it would not be able to account for year-round equipment and personnel contracting.

## 3. DESCRIPTION OF THE AFFECTED ENVIRONMENT AND ENVIRONMENTAL IMPACTS

### 3.1. INTRODUCTION

The discussion below will: (1) briefly describe/summarize the pertinent affected resources; (2) discuss whether proposed activities and their IPFs would have significant impacts to the human environment of the GOM; and (3) identify significant impacts, if any that would require further NEPA analysis in an EIS. The description of the affected environment and impact analysis are presented together in this section for each resource. For the impact analysis, resource-specific significance criteria were developed for each category of the affected environment. The criteria reflect consideration of both the context and intensity of the impact at issue (see 40 CFR § 1508.27). For the sake of this document, the criteria for impacts to environmental resources are classified into one of the three following levels:

- Significant Impact (including those that could be mitigated to non-significance);

- Adverse but Not Significant Impact; or
- Negligible Impact.

Preliminary screening for this assessment was based on a review of this relevant literature; previous SEAs, the Multisale EIS and 2018 SEIS, and statistics/data pertinent to historic and projected activities. BOEM initially considered the following resources for impact analysis:

- marine mammals (including ESA listed species);
- sea turtles (all are ESA listed species);
- fishes (including listed species and ichthyoplankton);
- commercial and recreational fisheries;
- coastal and marine birds (including ESA listed species);
- benthic communities (including deepwater benthic communities, live bottoms, and topographic features);
- archaeological resources;
- military uses;
- recreational and commercial diving;
- socioeconomic conditions (including employment, marine transportation, and infrastructure);
- geology/sediments; and
- air and water quality.

The impact analyses focus on a broad group of oil and gas activities and resources with the potential for non-negligible impacts. Routine, accidental, and cumulative impacts from exploration activities similar to those proposed by Anadarko are analyzed in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO that considered the proposed activities as well as impacts to resources relevant to the proposal. The level of impacts associated with each interaction was analyzed and described in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO and is incorporated by reference.

The Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO provides a comprehensive characterization of biological and socioeconomic resources that may be adversely affected by oil and gas exploration and development activities. For this SEA, BOEM evaluated the potential impacts resulting from the operator's proposed activities that were not considered in the Multisale EIS and 2018 SEIS. This section concentrates on the potential impacts of the proposed action on the following affected resources:

- air quality;
- offshore water quality;
- benthic biologically sensitive resources;
- marine mammals (including ESA listed species);
- sea turtles (all are ESA listed species);
- fish (including ESA listed species) and essential fish habitat (EFH); and
- archaeological resources.

Other environmental and socioeconomic conditions, identified in the initial list of resources considered for impact analysis above, such as military uses, were considered and the potential impacts that could occur from activities, such as the proposed activities, were fully addressed in the Multisale EIS and 2018 SEIS and deemed negligible (40 CFR § 1508.27) and are not discussed in this SEA. Space-use conflicts with recreational and commercial fishing vessels will be negligible compared to the area available for these activities, and there is a potential for an increase in some types of fishing activity due to development. There are no known recreational and/or commercial diving operations regularly occurring in the area. Although development could necessitate a negligible increase in commercial dive activity, potential impact levels do not warrant further analysis. Coastal and marine birds were not further analyzed due to the distance from shore and the temporary nature of the proposed activities. Topographic and pinnacle features were not further analyzed due to the distance from the proposed activities to the nearest topographic and/or pinnacle features (~82 mi and ~19 mi [~132 km and ~31 km] respectively). No socioeconomic effects were further analyzed due to the type, the temporary nature, and employment size, of the proposed activity. There is no expansion or modification of support bases proposed as a result of this activity. Additionally, support vessel operations are comparable to that described and analyzed in the Multisale EIS and 2018 SEIS for

similar activities. The potential impacts of a low-probability, Catastrophic Oil-Spill event, such as the *Deepwater Horizon* spill to the environmental resources and socioeconomic conditions listed above are fully addressed in the Catastrophic Spill Event Analysis (Appendix B of the 2012-2017 WPA/CPA Multisale EIS (USDOJ, BOEM, 2012). This analysis was later updated and published as a “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c) and a respective resource summary of that analysis is provided in each impact review below.

### ***Deepwater Horizon Impacts Incorporated into SEA Analyses***

BOEM, in conjunction with the well operator and other Federal and State agencies, continues to monitor and evaluate both the short-term and long-term impacts of the accidental spill. There is ongoing research to assess the impacts to resources from the *Deepwater Horizon* blowout, spill, and response efforts. For many resources, the data are still being collected and analyzed through the National Resource Damage Assessment (NRDA) process. BOEM continues to seek data and research results from the NRDA process and the scientific community. Results of this research are forthcoming, and BOEM subject matter experts are continuing to update their analyses as this information becomes available.

**Chapter 3** of this document describes the environmental and archaeological resources and the potential routine, accidental, and cumulative impacts of the proposed action on the resources that could be affected by the proposed activities. These descriptions present environmental resources as they are now, thus providing new baseline information that is informed by the *Deepwater Horizon* spill for analyses of potential impacts from the proposed activities.

## **3.2. AIR QUALITY**

### **3.2.1. Affected Environment**

The complete description of the air quality in the GOM region is set forth in Chapter 4.1 of the Multisale EIS and 2018 SEIS and is incorporated by reference. The following information is a summary of the description incorporated from the Multisale EIS and 2018 SEIS. Mississippi Canyon Blocks 80, 81, 82, 125, and 126 are located west of 87.5° W. longitude and falls under BOEM jurisdiction for enforcement of the Clean Air Act (CAA). The air over the OCS water is not classified, but some criteria pollutants may exceed the National Ambient Air Quality Standards (NAAQS) (USDOJ, BOEM, 2017a and b). The proposed exploration activities are located approximately 53 mi (85 km) from the nearest coastline of Plaquemines Parish, Louisiana. The Houston/Galveston, Texas area is in nonattainment for the 2008 ozone 8-hour standard. As of October 1, 2015, the United States Environmental Protection Agency (USEPA) strengthened NAAQS for ozone and will release new designated areas in the future after a comprehensive assessment of science, human health risk and exposure and alternative policy options before a rulemaking is proposed. Two new 1-hour NAAQS standards went into effect in 2010. They are the 1-hour NO<sub>2</sub> standard of 100 ppb and the 1-hour SO<sub>2</sub> standard of 75 ppb. The entire St. Bernard Parish, Louisiana is in nonattainment for the 2010 1-hr sulfur dioxide (SO<sub>2</sub>) standard, and part of the Evangeline Parish is in nonattainment for the same pollutant standard. Other than these areas, the coastal areas are in attainment of the NAAQS for ozone, carbon monoxide, nitrogen oxides, sulphur oxides, and particulate matter. For Prevention of Significant Deterioration (PSD) purposes, the coastal areas are classified as a Class II Areas.

Influences to onshore air quality are dependent upon meteorological conditions and air pollution emitted from operational activities. The pertinent meteorological conditions regarding air quality are the wind speed and direction, the atmospheric stability, and the mixing height (which govern the dispersion and transport of emissions). The typical, large-scale wind flow for the GOM area is driven by the clockwise circulation around the Bermuda High, resulting in a prevailing southeasterly to southerly wind flow, which is conducive to transporting air pollution emissions toward shore. However, superimposed upon this large-scale circulation are smaller scale wind-flow patterns, such as the land/sea breeze phenomenon. In addition, there are other large-scale weather features that occur periodically, namely tropical cyclones, and mid-latitude frontal systems. Because of the routine occurrence of these various conditions, the winds blow from all directions in the area of concern (MacDonald et al., 2004).

### 3.2.2. Impact Analysis

A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed activities on air quality can be found in Chapter 4.1 of the Multisale EIS and 2018 SEIS and is incorporated by reference. The following information is a summary of the impact analyses.

#### 3.2.2.1. Alternative 1

If selected, Alternative 1, the No Action Alternative, would result in the operator not undertaking the proposed activities as described in the plan. Therefore, site-specific IPFs to air quality would not occur. For example, there would be no volatile organic compound (VOC) emissions that would result in potential localized degradation of air quality. Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas related activity as described in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO and routine, accidental, and cumulative impacts would still occur from other activities.

#### 3.2.2.2. Alternative 2

If selected, Alternative 2, Proposed Action as Submitted and in accordance with their lease stipulations (in addition to any amendments), OCSLA requirements, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), and conditions of approval as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and FWS that concluded on April 20, 2018 (FWS 2018 BO). However, no additional, site-specific mitigation and monitoring measures would be required by BOEM or NMFS. As described in the analyses below, impacts to air quality from the proposed action are expected to be short-term, localized, and not lead to significant impacts.

#### Routine Activities

Air quality would be affected in the immediate vicinity of the proposed operations, service vessels, and aircraft. The impact from emissions for the proposed activities described in the Initial EP will not exceed BOEM's exemption levels per 30 CFR § 550.303(d), which would exempt the operator from additional air quality modeling. The proposed activities are not expected to significantly affect onshore air quality due to the distance from shore and the distance from the area of the proposed action to any PSD Class I air quality area, such as the Breton National Wildlife Refuge.

#### Accidental Events

Should a spill of oil occur, the VOCs, which would escape to the atmosphere from a surface slick, are precursors to photochemically produced ozone. A spike in VOCs could contribute to a corresponding spike in ozone, especially if the release were to occur on a hot sunny day in a NO<sub>2</sub>-rich environment. Should an accidental or emergency flaring or venting of gas occur, VOCs, methane, and carbon monoxide would also escape to the atmosphere. VOCs and nitrogen oxides can contribute to ozone formation in the presence of sunlight. Flared and vented gas can contain hydrogen sulfide which could result in emitted SO<sub>2</sub>. With the exception of Houston/Galveston, Texas area, the corresponding onshore area is in attainment for ozone. Additionally, with the exception of the St. Bernard Parish and part of the Evangeline Parish, Louisiana, the corresponding onshore area is in attainment for SO<sub>2</sub>. Due to the distance from shore, the proposed activities are not expected to have any impacts to onshore air quality, including nonattainment areas. If a fire occurs, prior to containment, particulate and combustible emissions will be released in addition to the VOCs. Emissions of pollutants into the atmosphere from routine activities associated with the proposed activities are expected to have minimal impacts to onshore air quality because of the prevailing atmospheric conditions, emission heights, emission rates, and the distance of these emissions from onshore.

Despite the recent *Deepwater Horizon* spill, historical trends in the GOM (see **Chapter 1.4** and the "Catastrophic Spill Event Analysis" white paper) indicate that catastrophic spill events are not likely to occur as a result of the activities associated with the proposed action. In the event of a catastrophic spill similar to the *Deepwater Horizon* spill, the "Catastrophic Spill Event Analysis" white paper (USDO, BOEM, 2017c) discusses the most likely and most significant impacts to air quality as it relates to the four phases of a major spill/blowout:

- 1) **Initial Event** (Section 2.2.1.1.; Page B-4);
- 2) **Offshore Spill** (Section 3.2.1.1; Page B-15);
- 3) **Onshore Contact** (Section 4.2.1.1; Page B-30); and
- 4) **Post-Spill, Long-Term Recovery** (Section 5.2.1.1.; Page B-40).

As the Catastrophic Spill Analysis in “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c) concludes, the potential impacts from a catastrophic spill could include air quality impacts that would require extensive recovery times.

## **Cumulative Impacts**

Cumulative impacts on air quality within the offshore area would come primarily from non-OCS oil/gas activities in the Gulf as well as sources on land such as generated outside the OCS and include emissions from industrial plants, power generation, and urban transportation. The location of the proposed action is far removed from coastal populations or industrial activity. The proposed activities are located over 53 mi (85 km) from the nearest shoreline, and would not affect the overall quality of air over the coast because of the distance to shore. Figure 4-1 of the Multisale EIS and 2018 SEIS (USDOJ, BOEM, 2017a and b) shows the Texas and Louisiana ozone attainment status. Except for Southeast Texas (Houston-Galveston-Brazoria), which is in nonattainment for ozone, and St. Bernard Parish and part of Evangeline Parish, Louisiana, which is in nonattainment for SO<sub>2</sub>, the Gulf’s coastal areas are currently designated as "attainment" for all of the NAAQS regulated pollutants (USEPA, 2020). Minor to moderate cumulative impacts on air quality are expected as a result of the proposed activities when added to the impacts of past, present, or reasonably foreseeable oil and gas development in the area, as well as other activities in the area, though the incremental impacts from the proposed activities are expected to be **minor**.

## **Conclusion**

The air quality in the immediate vicinity of the proposed activities would be affected by the projected emissions, but the 53 mi (85 km) distance between the area of the proposed action and the nearest shoreline results in substantial dilution factors for point-source emissions from the proposed action so that onshore air quality impacts would be well below levels considered to be significant.

### **3.2.2.3. Alternative 3**

If selected, Alternative 3, the Proposed Action with Additional Conditions of Approval, would allow the operator to undertake the proposed activities, as requested and conditioned in the plan and implement additional mitigation and monitoring measures as identified by BOEM and NMFS. As described in the analyses above for Alternative 2, impacts to air quality from the proposed action are expected to be localized and will not lead to significant impacts. The conditions of approval outlined in **Chapter 2.4** are not expected to increase or decrease the potential for effects to air quality from the proposed action. Alternative 3 does not differ from Alternative 2 because the additional mitigation and monitoring measures do not address this resource (i.e., all assumptions, estimates, and conclusions are identical); see the analysis provided in **Chapter 3.2** for Alternative 2 for this resource.

## **3.3. OFFSHORE WATER QUALITY**

### **3.3.1. Affected Environment**

The description of water quality in offshore waters of the GOM is set forth in Chapter 4.2 of the Multisale EIS and 2018 SEIS and is incorporated by reference. The following information is a summary of the description incorporated from the Multisale EIS and 2018 SEIS.

The GOM is the ninth largest waterbody in the world. The Mississippi River Basin drains 41 percent of the contiguous United States. The basin covers more than 1,245,000 square miles, and includes all or parts of 31 states and two Canadian provinces (USACE, 2020).

The physical oceanography of the deep Gulf can be approximated as a two-layer system with an upper layer about 800- to 1,000-m (2,625- to 3,281-ft) deep that is dominated by the Loop Current and associated clockwise (anticyclonic) eddies (Welsh et al., 2009; Inoue et al., 2008); and the lower layer below ~1,000 m (3,281 ft) that has near uniform currents (Welsh et al., 2009; Inoue et al., 2008).

Deep waters east of the Mississippi River are affected by the Loop Current and associated warm-core anticyclonic eddies, which consist of clear, low-nutrient water (Muller-Karger et al., 2001). Cold-core cyclonic eddies also form at the edge of the Loop Current and are associated with upwelling and nutrient-rich, high-productivity waters. More details on the physical oceanography of the GOM are available in Chapter 4.2 of the Multisale EIS and 2018 SEIS.

Typical water quality parameters that are considered important to the health of coastal and marine environments include temperature, salinity, dissolved oxygen, nutrients, pH, turbidity, and pollutants.

Surface water temperatures in the GOM vary seasonally from about 29 °C (84 °F) in the summer to about 19 °C (65 °F) in the winter (Gore, 1992). In the summer, warm water may be found from the surface down to a thermocline at depths to about 160 ft (50 m) deep. Minimum water temperatures at the deep seafloor approach 4° C (39 °F).

The salinity at the sea surface in the offshore central GOM is generally 36 parts per thousand (ppt) (Gore, 1992). Lower salinities are characteristic nearshore where fresh water from the rivers mix with shallow Gulf waters. For example, salinity in open water near the coast may vary between 29 and 32 ppt during fall and winter, but it may decline to 20 ppt during spring and summer due to increased runoff (USDOJ, MMS, 2000).

Dissolved oxygen (DO) concentrations in seawater vary as a function of temperature and barometric pressure. In general, cold water supports higher DO concentrations than warm water. DO concentrations between 5 and 10 milligrams per liter (mg/L) are considered beneficial to aquatic life. The GOM hypoxic zone is a band of oxygen-stratified water that stretches along the Texas-Louisiana shelf each summer where the DO concentrations are less than 2 mg/L. It is the largest hypoxic area in the entire western Atlantic Ocean (Turner et al., 2005). The hypoxic zone is the result of excess nutrients, primarily nitrogen, carried downstream by rivers to discharge to coastal waters. Density stratification results where the less dense, nutrient-rich freshwater spreads on top of the denser seawater and prevents oxygen from replenishing the bottom waters. The excess nutrients cause phytoplankton blooms which eventually die and sink to the bottom, where bacterial decomposition consumes DO.

Seawater generally averages pH 8 at the surface due to marine systems being buffered by carbonates and bicarbonates. However, in the open waters of the GOM, pH ranges from approximately 8.1 to 8.3 at the surface (Gore, 1992). The pH decreases to approximately 7.9 at a depth of 700 m (2,297 ft), and in deeper waters, it increases again to approximately 8.0 (Gore, 1992).

GOM coastal waters offshore of Texas, Louisiana, Mississippi, and Alabama exhibit high turbidity due to suspended sediment in river discharge, especially during seasonal periods of heavy precipitation. High turbidity may extend up to 50 mi offshore the Mississippi River and lesser distances to the east and west along the coast. Storms may also resuspend soft bottom sediments on the continental shelf, causing an increase in turbidity near the seafloor. Stratified water normally restricts this turbid water to within 20 m (66 ft) from the seafloor up into the water column (Bright et al., 1976; Bright and Rezak, 1978). Warm-core eddies can entrain and transport high turbidity shelf waters to farther offshore over deep Gulf waters. Outside of these areas, water clarity in the GOM is good to excellent, with low levels of suspended sediment.

River runoff may include pollutants such as nutrients, pesticides and other organic chemicals, and metals. The Mississippi River introduces approximately 3,680,938 bbl of oil and grease per year from land-based sources (NRC, 2003) into the waters of the GOM. Offshore waters, especially deeper waters, are more directly affected by natural seeps. Hydrocarbons enter the GOM through natural seeps at a rate of approximately 980,392 bbl per year (a range of approximately 560,224-1,400,560 bbl per year) (NRC, 2003).

The National Research Council estimated that, on average, approximately 26,324 bbl of oil per year entered Gulf waters from petrochemical and oil refinery industries in Louisiana and Texas. Spills to coastal waters include pipeline releases (annual estimate of 6,230 bbl), tank vessel incidents (5,390 bbl), and coastal facility releases (5,180 bbl); while spills to offshore waters include pipeline releases (annual estimate of 420 bbl) and tank vessel incidents (10,500 bbl) (NRC, 2003).

The April 2010 *Deepwater Horizon* oil spill resulted from failures of a cement well seal and subsea blowout preventer. The Government estimated that approximately 4.9 million barrels of oil were released during the event (Oil Spill Commission, 2011a), and that 1.84 million gallons of dispersant were used subsea at the wellhead and on the surface (Oil Spill Commission, 2011b). Additionally, the corresponding emission of methane from the wellhead during the event was estimated between  $9.14 \times 10^9$  and  $1.25 \times 10^{10}$

moles (Kessler et al., 2011). Short-term and long-term effects from the *Deepwater Horizon* oil spill are discussed in “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c).

### **3.3.2. Impact Analysis**

A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed activities on offshore water quality can be found in Chapter 4.2 of the Multisale EIS and 2018 SEIS, and is incorporated by reference. The IPFs associated with the proposed activities in Mississippi Canyon Blocks 80, 81, 82, 125, and 126 that could affect marine water quality include: (1) turbidity from bottom disturbances from well emplacement activities; (2) drilling discharges, including cuttings with associated drilling muds; and (3) accidental spills of crude oil, diesel fuel, chemicals, or other materials from vessels/blowouts in marine waters. As explained below, due to the type and the temporary nature of the proposed activities, no substantive impacts would be expected.

#### **3.3.2.1. Alternative 1**

If selected, Alternative 1, the No Action Alternative, would result in the operator not undertaking the proposed activities as described in the plan. Therefore, site-specific IPFs to offshore water quality would not occur. There would be no turbidity issues related to well emplacement activities that would result in potential localized degradation of water quality, no discharges during the drilling of the well and no accidental spills of crude oil, diesel fuel, chemicals, or other materials from vessels/blowouts in marine waters. Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas related activity as described in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO and routine, accidental, and cumulative impacts would still occur from other activities.

#### **3.3.2.2. Alternative 2**

If selected, Alternative 2, Proposed Action as Submitted and in accordance with their lease stipulations (in addition to any amendments), OCSLA requirements, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), and conditions of approval as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and FWS that concluded on April 20, 2018 (FWS 2018 BO). However, no additional, site-specific mitigation and monitoring measures would be required by BOEM or NMFS. As described in the analyses below, impacts to water quality from the proposed action, as submitted by the operator, are expected to be short-term, localized and not lead to significant impacts.

### **Routine Operations**

Impacts to water quality from routine activities associated with drilling or production may include overboard discharges of fluids and cuttings during drilling, development, and workovers of exploration and production wells; and service-vessel discharges.

The primary operational waste streams generated during offshore oil and gas exploration, development and production are drilling fluids, drill cuttings, various waters (e.g., bilge, ballast, fire, and cooling), deck drainage, sanitary wastes, domestic wastes, produced water, produced sand, and well treatment, workover, and completion fluids. Minor additional waste streams include desalination unit discharges, blowout preventer fluids, boiler blowdown discharges, excess cement slurry, several fluids used in subsea production, and uncontaminated freshwater and saltwater.

USEPA Regions 4 and 6 regulate all waste streams generated from offshore oil and gas activities. Section 403 of the Clean Water Act requires that National Pollutant Discharge Elimination System (NPDES) permits be issued for discharges to the territorial seas (baseline to 3 mi [5 km]), the contiguous zone, and the ocean in compliance with USEPA’s regulations for preventing unreasonable degradation of the receiving waters. Water Quality Standards consist of the waterbody’s designated uses, water quality criteria to protect those uses and to determine if they are being attained, and antidegradation policies to help protect high-quality waterbodies. Discharges from offshore activities near State water boundaries must comply with all applicable State Water Quality Standards. In general, waste streams that can be discharged overboard include water-based drilling fluids and drill cuttings, synthetic-based fluid-wetted drill cuttings,

cement slurries, various treated waters and sanitary wastes, and uncontaminated freshwater and saltwater provided they meet the criteria of the applicable NPDES permit.

Discharged water may not cause a sheen on the water surface, and the oil/grease concentration may not exceed 42 mg/L daily maximum, or 29 mg/L monthly average. The discharge must also be characterized for toxicity. The NPDES permits require no discharge within 1,000 m (3,281 ft) of an area of biological concern. Region 4 also requires no discharge within 1,000 m (3,281 ft) of any federally designated dredged material ocean disposal site.

Impacts to offshore waters from routine activities associated with the subject plan should be minimal. A detailed impact analysis of the routine impacts to offshore waters due to OCS activities can be found in Chapter 4.2 the Multisale EIS and 2018 SEIS.

## Accidental Events

Accidental events associated with the subject plan that could impact offshore water quality include spills of oil and refined hydrocarbons, releases of natural gas and condensate, spills of chemicals or drilling fluids, loss of well control, pipeline failures, collisions, or other malfunctions that would result in such spills. Spills from collisions are not expected to be significant. Overall, since major losses of well control and blowouts are rare events, potential impacts to offshore water quality are not expected to be significant except in the rare case of a catastrophic event. Although response efforts may decrease the amount of oil in the environment, the response efforts may also impact the environment through, for example, increased vessel traffic and the application of dispersants. Natural degradation processes will also decrease the amount of residual oil over time. Chemicals used in the oil and gas industry are not a significant risk to water quality because they are either nontoxic, are used in minor quantities, or are only used on a noncontinuous basis. A detailed impact analysis of the accidental impacts that may be associated with the proposed action on offshore waters can be found in Chapter 4.2 of the Multisale EIS and 2018 SEIS. Accidental spills as a result of a catastrophic event are discussed in “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c).

In the event of a catastrophic spill similar to the *Deepwater Horizon* spill, the “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c) discusses the most likely and most significant impacts to offshore water quality as it relates to three of the four phases of a major spill/blowout:

- 1) **Initial Event** (Section 2.2.1.2. ; Page B-5) ;
- 2) **Offshore Spill** (Section 3.2.1.2; Page B-16);
- 3) **Onshore Contact** (offshore water quality not included in this discussion); and
- 4) **Post-Spill, Long-Term Recovery** (Section 5.2.1.2.; Page B-40).

The potential impacts from a catastrophic spill could result in both temporary and long-term offshore water quality degradation that would require extensive recovery times. However, despite the *Deepwater Horizon* spill, historical trends in the GOM (see **Chapter 1.4** and the “Catastrophic Spill Event Analysis” white paper) indicate that catastrophic spill events are not likely to occur as a result of the proposed action.

## Cumulative Impacts

Exploration, development, and production activities contribute to cumulative water quality degradation in offshore waters. Surface spills of oil, diesel fuel, and other materials may occur from vessels transporting crude oil and petroleum products; from vessels involved in commercial fishing, freight or passenger transport; and from OCS operations. Such spills are low volume and are readily dispersed on the water surface. Well blowouts can disturb the bottom, increase turbidity, and put hydrocarbons into the sea. Should an oil spill  $\geq 1,000$  bbl (but not catastrophic) occur, localized, short-term changes in water quality would be expected; however, cumulative impacts on water quality over the long term would be **negligible**.

Therefore, no significant cumulative impacts on offshore water quality would be expected as a result of the proposed activities when added to the impacts of past, present, or reasonably foreseeable oil and gas development; as well as other activities in the area.

## Conclusion

Impacts on offshore water quality from the operational discharges that would be expected to result from the proposed action are negligible because of: 1) existing USEPA regulations; 2) water depth; 3) distance

of the project from the coast; 4) weathering; and 5) dilution factors. Spilled oil originating from the project is not expected to be  $\geq 1,000$  bbl and is expected to be substantially recovered/weathered while still at sea. Operator-initiated activities to contain and clean up an oil spill would begin as soon as possible after an event. Small quantities of unrecovered oil would weather and largely biodegrade within two weeks.

No significant long-term impacts on offshore water quality would be expected from the subject plan because of the type of and temporary nature of the proposed activity. Near-bottom water quality would be affected by increased turbidity and disturbed substrates during the period of well emplacement. Any effects from the elevated turbidity would be short term, localized, and reversible.

### **3.3.2.3. Alternative 3**

If selected, Alternative 3, the Proposed Action with Additional Conditions of Approval, would allow the operator undertaking the proposed activities, as requested and conditioned in the plan and implement additional mitigation and monitoring measures as identified by BOEM and NMFS. As described in the analyses above, impacts to offshore water quality from the proposed action are expected to be short-term, localized and not lead to significant impacts. The conditions of approval outlined in **Chapter 2.4** and discussed in the other resource sections are not expected to increase or decrease the potential for effects to offshore water quality from the proposed action. Alternative 3 does not differ from Alternative 2 because the additional mitigation and monitoring measures do not address this resource (i.e., all assumptions, estimates, and conclusions are identical); see the analysis provided in **Chapter 3.3** for Alternative 2 for this resource.

## **3.4. DEEPWATER BENTHIC COMMUNITIES**

For purposes of OCS activity impact analyses, BOEM defines “deepwater benthic communities,” to include chemosynthetic and deepwater coral communities in the GOM as those typically found in water depths of 984 ft (300 m) and greater (USDOJ, BOEM, 2017a and b).

### **3.4.1. Affected Environment**

A description of chemosynthetic and deepwater coral communities in the GOM region can be found in Chapter 4.4 of the Multisale EIS and 2018 SEIS. The following information is a summary of the descriptions in the EISs, and are incorporated by reference into this SEA.

The continental slope in the GOM extends from the edge of the continental shelf at a depth of about 656 ft (200 m) to a water depth of approximately 9,840 ft (3,000 m) (USDOJ, BOEM, 2017a and b). The vast majority of the GOM has a soft, muddy bottom in which burrowing infauna are the most abundant invertebrates. Mississippi Canyon Blocks 80, 81, 82, 125, and 126 fall into this category and the water depth of the proposed activities ranges from 3,710 to 4,305 ft (1,131 to 1,326 m).

A remarkable assemblage of invertebrates is found in association with hydrocarbon seeps in the GOM. Chemosynthetic communities can occur at or near hydrocarbon seeps and are defined as persistent, largely sessile assemblages of marine organisms dependent upon symbiotic chemosynthetic bacteria as their primary food source (MacDonald, 1992). Invertebrate taxa in these communities include tube worms and bivalves, among others. Symbiotic chemosynthetic bacteria live within specialized cells in the invertebrate organisms and are supplied with oxygen and chemosynthetic compounds (methane and sulfides) by the host via specialized blood chemistry (Fisher, 1990). Chemosynthetic bacteria, which live on mats, in sediment, and in symbiosis with chemosynthetic invertebrates, use a carbon source independent of photosynthesis to make sugars and amino acids. The host, in turn, lives off the organic products subsequently released by the chemosynthetic bacteria and may even feed on the bacteria themselves. Chemosynthetic communities can become established when a hard substrate is available for colonization at or near a seep. Depending on the situation, sessile benthic invertebrates can settle on and colonize carbonate substrate. These organisms form additional structure upon the seafloor, increasing the complexity of the habitat that may provide support to a variety of deepwater corals, invertebrates and fishes.

Some deepwater corals form communities occurring at or near hydrocarbon seeps, or on exposed outcrops, and may be found in association with chemosynthetic communities. Deepwater coral communities are also found on shipwrecks, and deepwater oil and gas infrastructure. These coral communities are distinctive and provide three-dimensional habitat for a range of fishes and invertebrates. Hard-bottom habitats in deep water include communities dominated by *Lophelia pertusa*, with other corals such as the bamboo coral (*Keratoisis flexibilis*) and zigzag coral (*Madrepora oculata*). Numerous other

invertebrates are also associated with these benthic habitats (Sulak et al., 2008; Cordes et al., 2008; Fisher et al., 2007; Schroeder et al., 2005).

Hydrocarbon seep communities in the GOM have been reported to occur at water depths greater than 300 m (984 ft) (USDOI, BOEM, 2017a and b). To date, there are over 300 documented deepwater benthic communities comprised of chemosynthetic organisms and/or deepwater corals. Once thought rare, research suggests that deepwater faunal communities are regularly associated with seafloor features commonly found in the vicinity of the primary geophysical signatures of the seabed for hydrocarbon migration to the seafloor. These areas include those where hydrocarbons percolate through sediments or where hydrocarbons move along faults that reach the seafloor. More than 23,000 positive anomalies have been identified from seismic survey data and each may represent a habitat where a hard substrate and a deepwater community may be found. However, until an anomaly has been visited and confirmed, it is unknown if hard substrates are exposed and capable of supporting deepwater benthic communities.

To map areas of probable habitat for deepwater benthic communities, scientists at BOEM analyzed decades of three-dimensional seismic data to classify seafloor returns exhibiting anomalously high or low reflectivity. The areas of high reflectivity represent patches of anomalous seafloor returns that likely indicate patches of hard seafloor that would provide substrate for deepwater benthic communities. Most confirmed hard bottoms in the deepwater GOM were created by the precipitation of calcium carbonate substrate by chemosynthetic bacterial activity and are capable of supporting deepwater benthic communities. However, non-biogenic hard bottoms are also found at escarpments, seafloor-reaching faults, or where salt formations reach the surface. Investigations of the seafloor at patches of high reflectivity indicate that chemosynthetic and coral communities are much more common in the deepwater GOM than previously known (USDOI, BOEM, 2017a and b). Also, areas of low reflectivity (negative anomalies) can be indicative of gassy sediments and mud volcanoes with a high flux of hydrocarbons from the seafloor. Although uncommon, chemosynthetic bivalves may be found in areas with a high flux of hydrocarbons.

### **3.4.2. Impact Analysis**

A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed activities on chemosynthetic communities and deepwater coral communities can be found in Chapter 4.4 of the Multisale EIS and 2018 SEIS. The following information is a summary of the impact analyses in the Multisale EIS and 2018 SEIS and it is incorporated by reference into this SEA.

Any hard substrate communities located in deep water would be particularly sensitive to impacts from OCS activities resulting in bottom disturbances and increased turbidity. Such impacts to these habitats could permanently prevent recolonization by similar organisms requiring hard substrate. The IPFs associated with the proposed activities in Mississippi Canyon Blocks 80, 81, 82, 125, and 126 that could affect deepwater benthic communities include physical impacts from: (1) well emplacement activities; (2) drilling discharges, including cuttings and drilling muds; and (3) seafloor blowouts without an oil spill during well drilling or emplacement of subsea infrastructure.

#### **3.4.2.1. Alternative 1**

If selected, Alternative 1, the No Action Alternative, would result in the operator not undertaking the proposed activities as described in the plan. Therefore, site-specific IPFs to deepwater benthic communities would not occur. For example, there would be no well emplacement activities that could result in physical damage to the deepwater benthic communities or their substrates, no drilling discharges that could result in burial of the organisms, or no burial due to a blowout. Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas related activity as described in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO and routine, accidental, and cumulative impacts would still occur from other activities.

#### **3.4.2.2. Alternative 2**

If selected, Alternative 2, the Proposed Action as Submitted and in accordance with their lease stipulations (in addition to any amendments), OCSLA requirements, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), and conditions of approval as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and FWS that concluded on April 20, 2018 (FWS 2018 BO). However, no additional,

site-specific mitigation and monitoring measures would be required by BOEM or NMFS. Examples of potential impacts to possible deepwater benthic communities include, but are not limited to, damage from well emplacement activities, smothering from drilling discharges, possible sedimentation and/or oil contamination from a blowout, and crushing or burial from emplacement of subsea infrastructure. Because the operator is required to follow all existing lease stipulations as well as the applicable regulations as clarified by NTLs (the operator reaffirmed compliance in its plan as cited above), conditions outlined in the following analyses related to NTL No. 2009-G40 will result in reducing the probability of impacts to deepwater benthic communities.

## **Routine Operations**

The NTL No. 2009-G40, (*Deepwater Benthic Communities*) provides guidance related to BOEM's regulations implementing a policy of avoidance of sensitive deepwater benthic communities or areas that have a high potential for supporting these community types, as interpreted from geophysical records. According to NTL No. 2009-G40, all plans submitted for deep water (984 ft, 300 m or greater) will be reviewed for the presence of deepwater benthic communities that may be affected by the proposed activity. Wells must be located a distance of at least 2,000 ft (610 m) from possible and known deepwater benthic communities to prevent cuttings from smothering the communities, and any seafloor disturbance (anchors, anchor chains, cables) must be at least 250 ft (76 m) from a possible or known deepwater benthic community. Lessees intending to explore or develop in water depths >984 ft (300 m) are required to provide information about geophysical surveys of the area of proposed activities and to evaluate the data for indications of conditions that may support sensitive deepwater benthic communities.

*Well Emplacement Activities:* Emplacement of the wells and associated subsea infrastructure can cause disturbances with lethal and sub-lethal effects such as (1) crushing; (2) burial; and (3) decreased fitness if substantial quantities of sediments are suspended in the water column during operations. For this plan, Anadarko proposes to use a DP drillship or DP semisubmersible to conduct their drilling activities; therefore, there are no anchors associated with the proposed operations. Also, the site-specific deepwater benthic communities review conducted for the proposed activities identified no areas or evidence of potential sensitive deepwater benthic communities or habitat that could support such communities within 2,000 ft (610 m) of the proposed well sites.

*Drilling Discharges:* Routine surface discharges from development drilling and production facility operations in water depths of 1,000 m (3,280 ft) can reach detectable accumulations at distances of at least 1 km (0.6 mi) (CSA, 2006); however, substantial sediment accumulations will be limited in distance from the surface discharge point. For discharges on the seafloor during initial well jetting, sediment accumulation may reach distances of approximately 100 m (328 ft) and could result in mounding in the immediate area around the well site (CSA, 2006). In both situations, splays of discharges tend to deposit in the direction of prevailing currents. Any discharges landing on any deepwater communities in substantial quantities during these activities could result in impacts directly due to mortality or indirectly due to sub-lethal impacts.

Distancing bottom disturbing activities from features that could support deepwater benthic communities, as described in NTL No. 2009-G40, minimizes potential impacts to deepwater benthic communities due to drilling discharges. Because of this distancing, any drill cuttings from deepwater operations would not come in contact with a deepwater community or would be diluted to such an extent to not result in negative impacts to a deepwater community. Because many deepwater organisms are long-lived with low reproductive rates, if a chemosynthetic community was impacted, it could take decades or centuries to recover depending on the size of the community. The deepwater benthic communities review conducted for this proposed action did not detect evidence indicating sensitive deepwater benthic communities or habitat that could support such communities within 2,000 ft (610 m) of the proposed well sites.

## **Accidental Events**

A blowout, as used here, is from expulsion of gas and/or water and/or suspended sediment and/or insubstantial oil out of a well. A blowout at the seafloor without the presence of substantial quantities of oil could occur when excess pressure in the well exceeds the capacity (both the operator's and the drilling apparatus' capacity) to contain the well. A blowout at the seafloor could create a crater on the sea bottom and/or suspend and disperse large quantities of bottom sediments, burying both infaunal (living in the sediment) and epifaunal (living on the sediment) organisms and interfering with sessile invertebrates that

rely on filter-feeding organs. Rapid burial by accumulations of sediment >1 ft (>30 cm) in thickness is likely to be lethal for all benthic organisms based on analysis of escape trace fossils from the geologic record (Frey, 1975; Basan et al., 1978; Eckdale et al., 1984). Lesser accumulations of sediment (or cuttings) may be lethal to some sessile (attached or immotile) invertebrates and survivable by motile organisms. Similar to impacts from drill cuttings, impacts from a blowout would be limited because of the duration and areal extent of the accident. Distancing the well at least 610 m (2,000 ft) from any feature that could support deepwater benthic communities also reduces that possibility of organisms being smothered by disturbed sediment. Any oil, sediments, or fluids released by a seafloor blowout of this nature could have potentially adverse effects on sensitive deepwater benthic communities. However, there are several reasons why substantive impacts from these are very unlikely for this IPF. First, the likelihood of any size blowout is very small. Since reporting requirements changed in 2006, there have been no reported blowouts of this nature (USDOJ, BSEE, 2020). Second, any sediments or fluids in this type of blowout would be limited in quantity, and the blowout would be limited in duration. As such, the sediments or fluids would either rise to the surface or be rapidly diluted in the water column and not impact deepwater communities given the proper distancing requirements. This type of blowout is not considered a catastrophic event similar to the *Deepwater Horizon* explosion and spill. For information on this type of event see the “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c).

## Cumulative Impacts

Operator adherence to lease stipulations will further ensure that the potential impacts of the proposed action on sensitive benthic communities and/or sites that could support them within the area of the proposed action will be low or non-existent, because there are no identified benthic communities near the proposed activities. Therefore, no significant cumulative impacts on sensitive benthic communities and/or sites that could support them would be expected and any impacts would be **negligible**.

## Conclusion

As mentioned earlier, a DP drillship or DP semi-submersible will be used for the proposed activities; therefore, anchors will not be required and the potential for bottom-disturbing activities will be reduced during routine activities. The site-specific benthic communities review conducted for the proposed action determined that there was no evidence of potentially sensitive benthic communities or sites that could support such communities within 2,000 ft (610 m) of the well site. Therefore, the impacts to deepwater benthic communities would be negligible.

### 3.4.2.3. Alternative 3

If selected, Alternative 3, Proposed Action with Additional Conditions of Approval, would allow the operator undertaking the proposed activities, as requested and conditioned in the plan, and implement additional mitigation and monitoring measures as identified by BOEM and NMFS. The conditions of approval outlined in **Chapter 2.4** and discussed in the other resource sections are not expected to increase or decrease the potential for effects to benthic resources from the proposed action. Alternative 3 does not differ from Alternative 2 because the additional mitigation and monitoring measures do not address this resource (i.e., all assumptions, estimates, and conclusions are identical); see the analysis provided in **Chapter 3.3** for Alternative 2 for this resource.

## 3.5. MARINE MAMMALS

### 3.5.1. Affected Environment

The marine mammal community is diverse and distributed throughout the northern GOM waters. The GOM’s marine mammals are represented by members of the taxonomic order Cetacea, including suborders Mysticeti (i.e., baleen whales) and Odontoceti (i.e., toothed whales), as well as the order Sirenia (i.e., manatee). Twenty-one species of cetaceans and one species of Sirenia regularly occur in the GOM and are identified in the NMFS Stock Assessment Reports (SAR) (Jefferson et al., 2008; Davis et al., 2000; Roberts et al., 2016; Hayes et al., 2018; Hayes et al., 2019; Hayes et al., 2020). A complete description of marine mammals can be found in Chapter 4.9 of the Multisale EIS and 2018 SEIS, and in the NMFS 2017, 2018, and 2019 SAR (Hayes et al., 2018, 2019, and 2020) and NMFS 2020 BO (USDOC, NMFS, 2020a), and are incorporated by reference.

## Threatened or Endangered Marine Mammal Species

Only two cetaceans, the sperm whale (*Physeter macrocephalus*) and the GOM Bryde's whale (*Balaenoptera edeni*), regularly occur in the GOM and are listed as endangered under the ESA. On January 8, 2016 (81 FR 999), the United States Fish and Wildlife Service (FWS) issued a proposed rule and notice to reclassify the West Indian manatee from endangered to threatened (*Federal Register*, 2016a) which was later issued as a Final Rule (82 FR 16668) on April 5, 2017 (*Federal Register*, 2017). On December 2, 1970, in the Final Rule (35 FR 18319), the sperm whale was listed as endangered throughout its range. The Final Rule (84 FR 15446) to list the GOM Bryde's whale as endangered was issued and became effective on May 15, 2019 (*Federal Register*, 2019). The only commonly occurring baleen whale in the northern GOM is the Bryde's whale. Most sightings have been made in the DeSoto Canyon region and off western Florida, although there have been some in the west-central portion of the northeastern GOM. The best estimate of abundance for Bryde's whales in the northern GOM is 33 individuals, which is the last estimate from a 2009 survey (Hayes et al., 2018 and 2020). Sperm whales in the GOM are not evenly distributed, showing greater densities in areas associated with oceanic features that provide the best foraging opportunities (USDOC, NMFS, 2020a).

## Non-ESA-Listed Marine Mammal Species

Nineteen toothed cetaceans (including beaked whales and dolphins) regularly occur in the GOM but are not ESA-listed. Despite being non-listed, the Marine Mammal Protection Act (MMPA) of 1972 protects all marine mammals.

## Unusual Mortality Events (UME)

An UME is defined under the MMPA as a “stranding that is unexpected, involves a significant die-off of any marine mammal population, and demands immediate response.” A list of active and closed UMEs with updated information can be found at the following website: <https://www.fisheries.noaa.gov/national/marine-life-distress/active-and-closed-unusual-mortality-events>.

### 2018-2020 Southwest Florida Bottlenose Dolphin UME (UME 66)

Since July 2018, the 2018-2020 Southwest Florida Bottlenose Dolphin UME was issued because of elevated bottlenose dolphin mortalities. Southwest Florida has been experiencing an ongoing severe bloom of a red tide organism since November 2017. The results from several necropsies showed positive findings of red tide toxin (brevetoxin) indicating this UME is related to the bloom. Other species such as fish, sea turtles, and manatees have also been impacted by the algal bloom (USDOC, NMFS, 2020b). This UME and its associated investigation are on-going.

## 3.5.2. Impact Analysis

A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed activities on marine mammals can be found in Chapters 4.9 of the Multisale EIS and 2018 SEIS, and is incorporated by reference. The IPFs with the proposed activities in Mississippi Canyon Blocks 80, 81, 82, 125, and 126 that could affect marine mammals include: (1) vessel noise and collisions; (2) marine debris; (3) water-quality degradation from drilling rig effluents; (4) oil spills and spill-response activities; and (5) drilling noise.

### 3.5.2.1. Alternative 1

If selected, Alternative 1, the No Action Alternative, would result in the operator not undertaking the proposed activities as described in the plan. Therefore, site-specific IPFs to marine mammals would not occur. Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas related activity as described in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO and routine, accidental, and cumulative impacts would still occur from other activities.

### 3.5.2.2. *Alternative 2*

If selected, Alternative 2, the Proposed Action as Submitted and in accordance with their lease stipulations (in addition to any amendments), OCSLA requirements, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), Reasonable and Prudent Measures and implementing Terms and Conditions as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and with FWS that concluded on April 20, 2018 (FWS 2018 BO). However, no additional, site-specific mitigation and monitoring measures would be required by BOEM or NMFS.

## **Routine Operations**

### *Vessel Traffic*

The proposed activities are expected to require several roundtrip supply-vessel and crew-vessel trips per week. Slow-moving cetaceans or those that spend extended periods of time at the surface might be expected to be the most vulnerable (Vanderlaan and Taggart, 2007). Smaller delphinids often approach vessels that are in transit to bow-ride; however, vessel strikes are less common for these faster moving mammals or are underreported (Wells and Scott, 1997). Florida manatees are commonly found in shallow coastal waters of Florida, but they have been found along the entire northern GOM from Florida to Texas (Fertl et al., 2005), though some deepwater sightings have occurred. Vessel strikes are the most common cause of human-induced mortality for manatees (State of Florida, Fish and Wildlife Conservation Commission, 2020), and most manatees bear prop scars from contact with vessels. The vast majority of strikes to manatees result from recreational and fishing vessels, not those related to oil and gas activities.

Worldwide, most vessel strikes of large whales occur when vessels are traveling at speeds greater than approximately 10 knots (11.5 miles per hour) (Conn and Silber, 2013; Jensen and Silber, 2004; Laist et al., 2001; Vanderlaan and Taggart, 2007). If a vessel strike occurs, the animal may experience no injuries, minor non-serious injuries, serious injuries, or death, which largely depends on the size and speed of the vessel (NMFS, 2020a). Both GOM Bryde's whales and sperm whales are vulnerable to vessel strikes. One confirmed vessel strike to a GOM Bryde's whale occurred in 2009. While there are no known recent vessel strikes to sperm whales, one possible lethal strike occurred in 1990 and a non-lethal strike in 2005. Additionally, a sperm whale is believed to have been struck by a U.S. Navy vessel in 2001 (USDOC, NMFS, 2020a).

The lack of response by sperm whales to oncoming vessels suggest the whales may not hear or see ships approaching or the whales are habituated to the high level of vessel operation activity in the GOM. The Bryde's whale spends much of its' time within 15 m (49 ft) of the water surface and at night on the surface, which makes it more likely to be struck by a vessel. With the Bryde's whale vessel strike mitigation measures required by the NMFS 2020 BO in place and as proposed under Alternative 2, NMFS estimated an annual rate of zero lethal Bryde's whale vessel strikes per year from oil and gas vessel traffic greater than 10 knots (USDOC, NMFS, 2020a). The proposed activities are located outside of the area where the Bryde's whale is likely to be present. The operator has not proposed any service vessels or vessel traffic within the Bryde's whale habitat area. Under Alternative 2, the operator is required to notify BOEM 15 days prior to any vessel transit within the Bryde's whale area; approval from BOEM is required prior to any vessel transit changes.

In their 2020 BO, NMFS estimated an annual rate of 0.10 vessel strikes likely to result in no or minor injuries to sperm whales per year from oil and gas activities (USDOC, NMFS, 2020a).

By selecting Alternative 2, the operator is required to follow mitigation and monitoring measures in Appendix C: Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols outlined in the NMFS 2020 BO. With these mitigation and monitoring measures in place, the impacts to the sperm whales, the Bryde's whale, and other marine mammals is determined to be **minor**.

### *Noise*

Marine mammals, including Bryde's whales and sperm whales, use sound in their environment to detect prey, predators, and habitat types, as well as for navigation and communication. Constant sounds in the environment can mask an animal's ability to communicate and hear important sounds within their environment (USDOC, NMFS, 2020a). In general, acute injury, or Permanent Threshold Shift (PTS), and disturbance would occur with the animal in close proximity to the sound. Temporary Threshold Shift (TTS)

is a short duration of auditory impairment that is usually recoverable within days or hours (USDOC, NMFS, 2020a).

The NMFS sets the 180-dB root-mean-squared (rms) isopleth where on-set of auditory injury or mortality (level A harassment) to cetaceans may occur. Southall et al. (2007) suggests this level should rather be at 230 dB rms for a nonpulsed sound, such as drilling noise. The source levels from drilling are relatively low (154 dB and below, as cited by Greene, 1986 in Richardson et al., 1995), below the level B (behavioral) harassment threshold of 160 dB set by NMFS under the MMPA. According to Southall et al. (2007), for behavioral responses to nonpulses (such as drill noise), data indicate considerable variability in received levels associated with behavioral responses. Contextual variables (such as novelty of the sound to the marine mammal and operation features of the sound source) appear to have been at least as important as exposure level in predicting response type and magnitude. There is little information on the behavioral responses by marine mammals to drilling noise in the GOM. Southall et al. (2007) summarized the existing research, stating that the probability of avoidance and other behavioral effects increases when received levels increase from 120 to 160 dB. Marine mammals may exhibit some avoidance behaviors, but their behavioral or physiological responses to noise associated with the proposed project are unlikely to have population-level impacts in the northern GOM.

### *Vessel Noise*

Vessel noise can have acute effects such as short-term behavioral and stress response. The nature of behavioral response cetaceans exhibit to vessels may depend on vessel speed, size, and distance from the animal, as well as the number and frequency of vessel encounters (USDOC, NMFS, 2020a). The dominant source of vessel sound from the proposed action is propeller cavitation, although other ancillary sounds may be produced. The intensity of sound from vessels is related to size and speed. Large ships tend to be noisier than small ones and ships underway with a full load or towing/pushing produce more sound than unladen vessels (USDOC, NMFS, 2020a). Noise from service-vessel traffic may elicit a startle and/or avoidance reaction from whales and dolphins or mask their sound reception. Vessel noise from the proposed action will produce low levels of noise, generally in the 150 to 170 dB re 1  $\mu$ Pa-m at frequencies below 1,000 Hz. Vessel noise is transitory and generally does not propagate at great distances from the vessel. The NMFS 2020 BO concluded that the effects of vessel noise to sperm whales are not likely to adversely affect the species and Bryde's whales are likely to be adversely affected from vessel noise (USDOC, NMFS, 2020a). The behavioral disruptions potentially caused by noise and the presence of service-vessel traffic will have **negligible** effects on cetacean populations in the northern GOM.

### *Aircraft Noise*

The noise and the shadow from helicopter overflights, take-offs, and landings can cause a startle response and can interrupt whales and dolphins while resting, feeding, breeding, or migrating (Richardson et al., 1995). The Federal Aviation Administration's Advisory Circular 91-36D (September 17, 2004) encourages pilots to maintain higher than minimum altitudes over noise-sensitive areas. Guidelines and regulations put in place by NOAA Fisheries under the authority of the MMPA include provisions specifying that helicopter pilots maintain an altitude of 1,000 ft (305 m) within 300 ft (91 m) of marine mammals. The proposed action is expected to have helicopter support with multiple transits between the MODU and airbase. The duration of the effects resulting from a startle response is expected to be short-term during routine flights, and the potential effects will be insignificant and not likely to adversely affect sperm and Bryde's whales (USDOC, NMFS, 2020a). Since these occurrences would be temporary and pass within seconds, marine mammals are not expected to be adversely affected by routine helicopter traffic operating at prescribed altitudes and impacts are considered **negligible**.

### *Drilling and Production Noise*

Offshore drilling and production involve a variety of activities that produce underwater sounds at intensities and frequencies that can be heard by cetaceans. The sound sources from fixed or stationary production platforms are localized and are not expected to produce sound levels over great distances. Noise from drilling could be continuous or intermittent, sudden, and at times high intensity as operations take place. Sounds emanating from drilling activities from fixed platforms are considered not very intense and generally are at very low frequencies near five Hz (USDOC, NMFS, 2020a). Gales (1982) reported received levels of 119 to 127 dB re 1  $\mu$ Pa-m at near-field measurements, while other measurements have recorded higher levels of sound up to 185 dB (rms) from platforms or 195 dB (rms) from drill ships. The

estimated frequencies of drilling from semisubmersible vessels are broadband from 80-4,000 Hz with an estimated source level (SL) of 154 dB re 1 $\mu$ Pa at 1 m. The tone of 60 Hz were SLs of 149 dB, 181 Hz were 137 dB, and 301 Hz were 136 dB (Greene, 1986). Drillships show somewhat higher sound levels as a result of mechanical sounds generated through the drillship hull and by the use of thrusters to maintain position while drilling. Sounds from semisubmersible platforms also show rather low sound source levels. In general, the sounds associated with offshore oil and gas exploration and production are generally at low levels and typically at very low frequencies (~4.5 to 38 Hz) (Gales, 1982).

### ***Casing Conductor (Drive Pipe) Installation***

Due to the frequency of exploratory and development drilling operations on the OCS, the greatest number of pile-driving operations involve the setting or installation of casing conductors. Most casing conductors range in diameter from 12-36 inches and have wall thicknesses that run from 0.25-0.75 inches. These are generally driven into the substrate until the conductor “meets refusal” or cannot be driven further without damage. Conductor casings can also be jetted into the seabed; however, the ease of mobilization of hammer drivers coupled with their speed of penetration, minimizes the use of jetting equipment, which requires more time to deploy and is often unviable due to water depth and sediment type. Most casing conductor driving operations occur in water depths less than 200 m (656 ft).

### ***Dynamically Positioned Vessels***

The potential effects that water-transmitted noise has on marine mammals include disturbance (subtle changes in behavior, interruption of previous activities, or short- or long-term displacement), masking of sounds (calls from conspecifics, reverberations from own calls, and other natural sounds such as surf or predators), physiological stress, and hearing impairment. Individual marine mammals exposed to recurring disturbance could be negatively affected. The temporary and transient noise associated with drilling and production is not expected to produce more than negligible to minor impacts on marine mammals since they are not expected in amplitudes sufficient to cause hearing or behavioral effects, and due to the wide-ranging behavior of marine mammal species. NMFS determined stationary and localized effects of platform-associated sounds, sperm whale encounters near platforms would be very brief as they swim by, and the potential effects of these sounds to disturb sperm whales will be insignificant. Construction and operation sounds other than pile driving will have insignificant effects on Bryde’s whales.

### ***Marine Debris***

Marine debris has the potential to adversely affect marine mammals through ingestion and entanglement. These effects could result in reduced feeding, reduced reproductive success, and potential injury, infection, or death. NMFS concluded marine debris is likely to adversely affect sperm and Bryde’s whales. Implementation of Appendix B: Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols and the International Convention for the Prevention of Pollution from Ships (MARPOL) requirements under the United States Coast Guard (USCG) may reduce, but not eliminate the risk of marine debris (USDOC, NMFS, 2020a). Without implementation of mitigations, marine mammals would be more vulnerable to direct impacts from entanglement in or ingestion of OCS marine debris. The impacts from OCS marine debris with implementation of mitigations are expected to be **negligible**.

### ***Water Degradation***

Waste streams generated by the drilling process include drilling fluids, drill cuttings, produced water, and deck drainage. The platform also contributes sanitary and domestic wastes. During production, there are additional waste streams including produced sand and well treatment, workover, and completion fluids. Further minor discharges include releases from desalination units, blowout preventer fluids, boiler blowdown, and excess cement slurry.

The USEPA regulates discharges from oil and gas operations to offshore marine waters in the GOM through NPDES general permits. Discharges of produced water, drilling fluids, drill cuttings, and chemically treated miscellaneous discharges under the NPDES general permit will be required to meet the whole effluent toxicity requirements. NMFS determined that because of USEPA regulation, most of the routinely discharged chemicals are not expected to result in exposure intensities that would adversely affect listed species because they are diluted and dispersed when released in marine waters. Without implementation of discharge requirements under the NPDES permit, marine mammals would be more

vulnerable to direct impacts from degraded water quality; therefore, potential impacts to marine mammals from water quality impacts are expected to be **negligible**.

## Accidental Events

### *Oil Spills and Response Activities*

BOEM defines a very large spill, as any spill volume greater than or equal to 10,000 bbl, and provided NMFS with information projecting that two oil spills greater than or equal to 10,000 bbl may occur over the duration of 10 years (through 2029); however, a defined upper volume for such a spill size cannot be predicted. BOEM “does not consider an extremely large event as reasonably certain to occur” over the time frame of the NMFS 2020 BO, although BOEM does acknowledge that impacts from the *Deepwater Horizon* spill warranted inclusion in NMFS 2020 consultation and BO as part of the environmental baseline.

The oil from an oil spill can adversely affect cetaceans by causing soft tissue irritation, fouling of baleen plates, respiratory stress from inhalation of toxic fumes, food reduction or contamination, direct ingestion of oil and/or tar, and temporary displacement from preferred habitats. The long-term impacts to marine mammal populations are poorly understood but could include decreased survival and lowered reproductive success. The range of toxicity and degree of sensitivity to oil hydrocarbons and the effects of cleanup activities on cetaceans are unknown. One notion concerning the use of dispersants is that chemical dispersion of oil will considerably reduce the impacts to aquatic mammals, primarily by reducing their exposure to petroleum hydrocarbons (French-McCay 2004; NRC, 2005). Dispersants are chemicals that reduce surface tension between oil and water, leading to oil droplet formation, so that the oil will more readily disperse into the water column. Dispersants typically contain surfactants and solvents and are used to entrain oil in the water column so as to protect shorelines from floating oil, but in turn, increases exposure to underwater organisms (USDOC, NMFS, 2020a). Chemical dispersant application during an oil spill may lower the amount of oil to which an aquatic mammal is exposed while increasing the potential loss of the insulative properties of fur through the reduction of surface tension at the fur-water interface (NRC, 2005).

Impacts from the dispersants are unknown but dispersants may contain ingredients that are known to irritate sensitive tissues of marine mammals (NRC, 2005). There have been no experimental studies and only a handful of observations suggesting that oil has harmed any manatees (St. Aubin and Lounsbury, 1990). Types of impacts to manatees from contact with oil include: (1) asphyxiation due to inhalation of hydrocarbons; (2) acute poisoning due to contact with fresh oil; (3) lowering of tolerance to other stressors due to the incorporation of sublethal amounts of petroleum components into body tissues; (4) nutritional stress through damage to food sources; and (5) inflammation or infection and difficulty eating due to oil sticking to the sensory hairs around their mouths (Preen, 1989, in Sadiq and McCain, 1993, AMSA, 2003). For a population whose environment is already under great pressure, even a localized incident could have population level consequences (St. Aubin and Lounsbury, 1990). Spilled oil might affect the quality or availability of aquatic vegetation, including seagrasses, upon which manatees feed.

In the event of catastrophic spill similar to the *Deepwater Horizon* spill, the “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c) discusses the most likely and most significant impacts to marine mammals as it relates to the four phases of a major spill/blowout:

- 1) **Initial Event** (Section 2.2.2.3. ; Page B-6) ;
- 2) **Offshore Spill** (Section 3.2.2.3; Page B-18);
- 3) **Onshore Contact** (Section 4.2.2.3; Page B-32); and
- 4) **Post-Spill, Long-Term Recovery** (Section 5.2.2.3; Page B-41).

More detailed information can be found in the Multisale EIS and 2018 SEIS.

## Conclusion

The sections above discuss the potential range of effects to marine mammals from the proposed activity and any of these effects has the potential individually or cumulatively to result in impacts to marine mammal species commonly found in the GOM and proposed action area. However, BOEM finds that the potential for such effects from the proposed action are unlikely to rise to significant levels for the following reasons:

- Mysticetes, as low-frequency hearing specialists, are the species groups most likely to be susceptible to impacts from nonpulse sound (intermittent or continuous) given that their hearing ranges overlap

most closely with the noise frequencies produced from drilling (Southall et al., 2007). However, most mysticete species that may occur in the GOM (i.e., North Atlantic right, blue, fin, sei, humpback, and minke) are considered either “extralimital,” “rare,” or “uncommon” within the GOM (Wursig et al., 2000; Hayes et al., 2019 and 2020). The only commonly occurring baleen whale in the northern GOM is the Bryde’s whale which is limited in its range. Given the small geographic scope of the proposed action, the presence of these species within the action area is not anticipated.

- The remaining marine mammal species in the GOM are considered either mid-frequency hearing specialists (e.g., sperm whales, beaked whales, and dolphins) with hearing ranges that slightly overlap with sound frequencies produced from drilling noise (Southall et al., 2007), or high-frequency specialists (pygmy and dwarf sperm whales). It is expected that there will be some overlap in the frequencies of the drill source and the hearing thresholds of the marine mammals present in the GOM. Wartzok and Ketten (1999) stated that bottlenose dolphins have hearing thresholds ranging from less than 5 kHz to over 100 kHz. Ridgway and Carder (2001) found, through auditory brainstem analysis, that pygmy sperm whales have thresholds from 90 to 150 kHz. Gordon et al. (1996) found that a stranded sperm whale had lower hearing limits at around 100 Hz while Ridgway and Carder (2001) found that a sperm whale calf had best hearing sensitivity between 5 and 20 kHz. Since there is some overlap in drilling and vessel sound levels produced and hearing thresholds of marine mammals, there is potential for the drilling noise produced to cause auditory and non-auditory effects, PTS, TTS, behavioral changes, or masking but it is expected to be limited.
- The estimated source levels of drilling from semisubmersible vessels (Greene, 1986) all fall below the 180 dB Level A harassment isopleths.
- The operator proposes adherence with the guidance provided under the NMFS 2020 BO Appendix B: Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols, which appreciably reduces the likelihood of marine mammals encountering marine debris from the proposed activity.

The geographic scope of the proposed action is small in relation to the ranges of marine mammals in the GOM. The proposed activities are not expected to cause long-term or permanent displacement of the animals from preferred habitats, nor will they result in the destruction or adverse modification of any habitats. In conclusion, because of the scope, timing, and transitory nature of the proposed action and the condition(s) of approval and monitoring requirements in place, the noise related to the proposed drilling operation is not expected to result in PTS, TTS, behavioral change, masking, or non-auditory effects to marine mammals in the GOM that would rise to the population level.

## Cumulative Impact Analysis

The proposed action may cumulatively affect cetaceans when viewed in light of the *Deepwater Horizon* explosion, spill, and response. Oil and gas leasing, exploration, development and production activities could impact marine mammals from the degradation of water quality resulting from operational discharges; vessel traffic; noise generated by platforms, drilling rigs, helicopters, and vessels; seismic surveys; explosive structure removals; oil spills; oil-spill-response activities; and loss of debris from service vessels and OCS structures. The cumulative impact on marine mammals is expected to result in a number of chronic and sporadic sublethal effects (i.e., behavioral effects and nonfatal exposure to or intake of OCS-related contaminants or discarded debris) that may stress and/or weaken individuals of a local group or population and predispose them to infection from natural or anthropogenic sources (Harvey and Dahlheim, 1994).

Few deaths are expected from chance vessel collisions and ingestion of plastic material. Disturbance (noise from vessel traffic and drilling operations, etc.) and/or exposure to sublethal levels of toxins and anthropogenic contaminants may stress animals, weaken their immune systems, and make them more vulnerable to parasites and diseases that normally would not be fatal (Harvey and Dahlheim, 1994). The net result of any disturbance will depend upon the size and percentage of the population likely to be affected, the ecological importance of the disturbed area, the environmental and biological parameters that influence an animal’s sensitivity to disturbance and stress, and the accommodation time in response to prolonged disturbance (Geraci and St. Aubin, 1980).

The effects of the proposed action, when viewed in light of the effects associated with other relevant activities, may impact marine mammals in the GOM. However, the operator is required to follow all existing lease stipulations, regulations, NTLs, and mitigation and monitoring measures provided in the

NMFS 2020 BO under Appendix B: Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols; Appendix C: Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols; and Appendix E: Summary of Oil Industry Discharges to the OCS Authorized by USEPA General NPDES Permits. Because of the operator's reaffirmed compliance with regulatory requirements, as well as the limited scope, timing, and geographic extent of the proposed action, effects from the proposed activities on marine mammals will be **negligible**. Therefore, no population level cumulative impacts to marine mammals would be expected as a result of the proposed activities when added to the impacts of past, present, or reasonably foreseeable oil and gas development in the area as well as other activities in the area. More detailed information can be found in the Multisale EIS and 2018 SEIS.

### **3.5.2.3. Alternative 3**

If selected, Alternative 3, the Proposed Action with Additional Conditions of Approval, would allow the operator undertaking the proposed activities, as requested and conditioned in the plan; however, the operator would be required to undertake additional mitigation and monitoring measures as identified by BOEM and NMFS. Alternative 3 differs from Alternative 2 because the additional mitigation and monitoring measures address this resource. NMFS provided concurrence, in that, the proposed activities and the use of a moon pool are within the scope of the effect analysis of the NMFS 2020 BO. Because the operator is required to follow all existing lease stipulations and regulations as clarified by NTLs, conditions outlined in the previous analyses related to BOEM NTL Nos. 2015-G03, 2016-G01, and 2016-G02; and the requirement for additional mitigation and monitoring measures should minimize or reduce the risk of a significant impact to marine mammals.

## **3.6. SEA TURTLES**

### **3.6.1. Affected Environment**

The life history, population dynamics, status, distribution, behavior, and habitat use of sea turtles can be found in Chapter 4.9 of the Multisale EIS and 2018 SEIS, and is incorporated by reference. Of the extant species of sea turtles, five are known to inhabit the waters of the GOM (Pritchard, 1997): the leatherback, green, hawksbill, Kemp's ridley, and loggerhead. These five species are all highly migratory, and individual animals will migrate into nearshore waters as well as other areas of the North Atlantic Ocean, GOM, and Caribbean Sea. All five species of sea turtles found in the GOM have been federally listed as endangered or threatened since the 1970's. Critical habitat has been designated for the Northwest Atlantic Ocean Loggerhead sea turtle distinct population segment (DPS) in the GOM (*Federal Register*, 2014).

In 2007, FWS and NMFS published 5-year status reviews for all federally listed sea turtles in the GOM (USDOC, NMFS and USDO, FWS, 2007a-e). A 5-year review is an ESA-mandated process that is conducted to ensure that the listing classification of a species as either threatened or endangered is still accurate. As of 2013, two 5-year reviews have been updated for the Leatherback and Hawksbill sea turtles (USDOC, NMFS and USDO, FWS, 2013a and b). Both agencies share jurisdiction for federally listed sea turtles and jointly conducted the reviews. After reviewing the best scientific and commercially available information and data, the agencies determined that the current listing classification for the five sea turtle species remain unchanged.

### **3.6.2. Impact Analysis**

A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed exploration activities on sea turtles can be found in Chapter 4.9 of the Multisale EIS and 2018 SEIS, and is incorporated by reference. The diversity of a sea turtle's life history leaves it susceptible to many natural and human impacts, including impacts while it is on land, in the benthic environment, and in the pelagic environment. The IPFs associated with the proposed activities in Mississippi Canyon Blocks 80, 81, 82, 125, and 126 that could affect sea turtles include: (1) vessel noise and collisions; (2) marine debris; (3) water-quality degradation from drilling rig effluents; (4) oil spills and spill-response activities; and (5) drilling noise.

#### **3.6.2.1. Alternative 1**

If selected, Alternative 1, the No Action Alternative, would result in the operator not undertaking the proposed activities as described in the plan. Therefore, site-specific IPFs to sea turtles would not occur.

Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas related activity as described in the Multisale EIS, 2018 SEIS, FWS 2018 BO, and NMFS 2020 BO and routine, accidental, and cumulative impacts would still occur from other activities.

### **3.6.2.2. Alternative 2**

If selected, Alternative 2, the Proposed Action as Submitted and in accordance with their lease stipulations (in addition to any amendments), OCSLA requirements, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), Reasonable and Prudent Measures and implementing Terms and Conditions as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and with FWS that concluded April 20, 2018 (FWS 2018 BO). However, no additional, site-specific mitigation and monitoring measures would be required by BOEM or NMFS.

## **Routine Operations**

### ***Vessel Noise***

The dominant source of noise from vessels is propeller operation, and the intensity of this noise is largely related to ship size and speed. Vessel noise from the proposed action would produce low levels of noise, generally in the 150 to 170 dB re 1  $\mu$ Pa-m at frequencies below 1,000 Hz. Vessel noise is transitory and generally does not propagate at great distances from the vessel. Also, available information indicates that sea turtles do not greatly utilize environmental sound. The NMFS 2020 BO concluded that sound sources associated with vessel movement were not likely to adversely affect sea turtles (USDOC, NMFS 2020a).

Drilling activities would produce sounds transmitted into the water that could be intermittent, sudden, and at times could be high intensity as operations take place. However, sea turtles are not expected to be impacted by this disturbance.

Popper et al. (2014) published sound exposure guidelines for fishes and sea turtles. The guidelines were broad-ranging and provided non-quantified, generalized guidelines for shipping noise as a low risk of impairment, unless the turtle is in the near field range (within tens of meters), which would pose a moderate risk of TTS that can recover over time. The risk for noise to cause masking and behavior effects range from low to high depending on the location of the turtle relative to the noise (Popper et al., 2014).

### ***Vessel Collisions***

Sea turtles spend at least 3-6 percent of their time at the surface for respiration and perhaps as much as 26 percent of time at the surface for basking, feeding, orientation, and mating (Lutcavage et al., 1997). Based on the behavioral observations of sea turtle avoidance of small vessels, green turtles may be susceptible to vessel strikes at speeds as low as two knots (2.3 mph) (Hazel et al. 2007). Although there have been hundreds of thousands of vessel trips that have been made in support of offshore operations during the past 40 years of OCS oil and gas operations, there have been no reports of OCS-related vessels having struck sea turtles. This is most likely because a strike with a sea turtle would probably go undetected by larger vessels and strikes are not reported. Despite the lack of on-water reporting, stranding records show that interactions between vessels and sea turtles in the GOM are common (USDOC, NMFS, 2020a). Data show that collisions with all types of commercial and recreational vessels are a cause of sea turtle mortality in the GOM (Lutcavage et al., 1997). Stranding data for the U.S. Gulf and Atlantic Coasts, Puerto Rico, and the U.S. Virgin Islands show that between 1986 and 1993 about 9 percent of living and dead stranded sea turtles had boat strike injuries (Lutcavage et al., 1997). Vessel-related injuries were noted in 13 percent of stranded turtles examined from the GOM and the Atlantic during 1993 (Teas, 1994), but this figure includes those that may have been struck by boats post-mortem. Large numbers of loggerheads and 5-50 Kemp's ridley turtles are estimated to be killed by vessel traffic per year in the U.S. (NRC, 1990; Lutcavage et al., 1997).

There have been no known documented sea turtle collisions with drilling and service vessels in the GOM; however, collisions with small or submerged sea turtles may go undetected. Based on sea turtle density estimates in the GOM, the encounter rates between sea turtles and vessels would be expected to be greater in water depths less than 200 m (USDOC, NMFS, 2007). Additionally, recent satellite tracking

studies have provided data to support that larger turtles often remain closer to shore to feed, nest and/or migrate; for loggerheads (Hart et al., 2013 and 2014) and Kemp's ridleys (Shaver et al., 2014). By selecting Alternative 2, the operator is required to follow mitigation and monitoring measures in Appendix C: Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols outlined in the NMFS 2020 BO. With these mitigation and monitoring measures in place, the impacts to sea turtles is determined to be **minor**.

With implementation of these measures and the avoidance of potential strikes from OCS vessels, the NMFS 2020 BO concluded that the risk of collisions between oil/gas-related vessels (including those for geological and geophysical [G&G] surveys, drilling, production, decommissioning, and transport) and sea turtles is appreciably reduced, but strikes may still occur. BOEM monitors for any takes that have occurred as a result of vessel strikes and also requires that any operator immediately report the striking of any animal.

To date, there have been no known or reported strikes of sea turtles by drilling vessels. Given the scope, timing, and transitory nature of the proposed action and with this established condition(s) of approval, effects to sea turtles from drilling vessel collisions is expected to be **negligible**.

### ***Marine Debris***

Many types of plastic materials end up as solid waste during drilling and production operations. Some of this material is accidentally lost overboard where sea turtles could consume it or become entangled in it. The incidental ingestion of marine debris and entanglement could adversely affect sea turtles. Marine debris is a continuing problem for sea turtles. Sea turtles living in the pelagic environment commonly eat or become entangled in marine debris (e.g. tar balls, plastic bags/pellets, balloons, and ghost fishing gear) as they feed along oceanographic fronts where debris and their natural food items converge. This is especially problematic for sea turtles that spend all or significant portions of their life cycle in the pelagic environment (i.e., leatherbacks, juvenile loggerheads, and juvenile green turtles) (USDOC, NMFS, 2020a). Implementation of Appendix B: Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols and the International Convention for the Prevention of Pollution from Ships (MARPOL) requirements under the United States Coast Guard (USCG) may reduce, but not eliminate the risk of marine debris (USDOC, NMFS, 2020a). Without implementation of mitigations, sea turtles would be more vulnerable to direct impacts from entanglement in or ingestion of OCS marine debris. The impacts from OCS marine debris with implementation of mitigations are expected to be **negligible**.

### ***Water Degradation***

Most operational discharges are diluted and dispersed when released in offshore areas and are considered to have sublethal effects (NRC, 1983; API, 1989; Kennicutt, 1995; Kennicutt et al., 1996). Any potential impacts from drilling fluids would be indirect, either as a result of impacts to prey species or possibly through ingestion via the food chain (Neff et al., 1989). The USEPA regulates discharges from oil and gas operations to offshore marine waters in the GOM through NPDES general permits. Discharges of produced water, drilling fluids, drill cuttings, and chemically treated miscellaneous discharges under the NPDES general permit will be required to meet the whole effluent toxicity requirements. NMFS determined that because of USEPA regulation, most of the routinely discharged chemicals are not expected to result in exposure intensities that would adversely affect listed species being that they are diluted and dispersed when released in marine waters. Without implementation of discharge requirements under the NPDES permit, sea turtles would be more vulnerable to direct impacts from degraded water quality. Impacts from water degradation are expected to be negligible due to the localized nature of the proposed activity and the wide-ranging habits of sea turtle species in the GOM.

## **Accidental Events**

### ***Oil Spills and Response Activities***

The oil from an oil spill can adversely affect sea turtles by causing soft tissue irritation, respiratory stress from inhalation of toxic fumes, food reduction or contamination, direct ingestion of oil and/or tar, and temporary displacement from preferred habitats (Lutz and Lutcavage, 1989). The long-term impacts to sea turtle populations are poorly understood but could include decreased survival and lowered reproductive success. The range of toxicity and degree of sensitivity to oil hydrocarbons and the effects of cleanup activities on sea turtles are unknown. Impacts from the dispersants are unknown, but may have

similar irritants to tissues and sensitive membranes as they are known to have had on seabirds and marine mammals (NRC, 2005).

In the event of a catastrophic spill similar to the *Deepwater Horizon* spill, the “Catastrophic Spill Event Analysis” white paper (USDOJ, BOEM, 2017c) discusses the most likely and most significant impacts to sea turtles as it relates to the four phases of a major spill/blowout:

- 1) **Initial Event** (Section 2.2.2.4. ; Page B-7) ;
- 2) **Offshore Spill** (Section 3.2.2.4; Page B-19);
- 3) **Onshore Contact** (Section 4.2.2.4; Page B-33); and
- 4) **Post-Spill, Long-Term Recovery** (Section 5.2.2.4; Page B-41).

In the event of a catastrophic spill similar to the *Deepwater Horizon* spill, any substantive impact to sea turtles is very unlikely because the potential impacts from a catastrophic spill would be similar to the aforementioned routine and accidental issues. However, despite the *Deepwater Horizon* spill, historical trends in the GOM (see **Chapter 1.4**) indicate that catastrophic spill events are not likely to occur as a result of the activities associated with the proposed action.

### **Cumulative Impact Analysis**

Activities considered under the cumulative scenario, including the proposed action, may affect sea turtles. Sea turtles may be impacted by oil and gas leasing, exploration, development and production activities including the degradation of water quality resulting from operational discharges, vessel traffic, noise generated by platforms, drilling rigs, helicopters and vessels, seismic surveys, explosive structure removals, oil spills, oil-spill-response activities, loss of debris from service vessels and OCS structures, commercial fishing, capture and removal, and pathogens. The cumulative impact of these ongoing OCS activities on sea turtles is expected to result in a number of chronic and sporadic sublethal effects (i.e., behavioral effects and nonfatal exposure to or intake of OCS-related contaminants or discarded debris) that may stress and/or weaken individuals of a local group or population and that may predispose them to infection from natural or anthropogenic sources.

Few deaths are expected from chance collisions with OCS service vessels, ingestion of plastic material, commercial fishing, and pathogens. Disturbance (noise from vessel traffic and drilling operations, etc.) and/or exposure to sublethal levels of toxins and anthropogenic contaminants may stress animals, weaken their immune systems, and make them more vulnerable to parasites and diseases that normally would not be fatal during their life cycle. The net result of any disturbance depends upon the size and percentage of the population likely to be affected, the ecological importance of the disturbed area, the environmental and biological parameters that influence an animal’s sensitivity to disturbance and stress, or the accommodation time in response to prolonged disturbance (Geraci and St. Aubin, 1980). As discussed above, lease stipulations and regulations are in place to reduce vessel strike mortalities.

Incremental injury effects from the proposed action on sea turtles are expected to be **negligible** for drilling and vessel noise, and **minor** for vessel collisions, but will not rise to the level of significance because of the limited scope, duration, and geographic area of the proposed drilling and vessel activities and the relevant regulatory requirements.

The effects of the proposed action, when viewed in light of the effects associated with other relevant activities, may affect sea turtles occurring in the GOM. With the enforcement of regulatory requirements for drilling and vessel operations and the scope of the proposed action, incremental effects from the proposed activities on sea turtles will be **negligible** (drilling and vessel noise) to **minor** (vessel strikes). The best available scientific information indicates that sea turtles do not greatly use sound in the environment for survival; therefore, disruptions in environmental sound would have little effect. Consequently, no significant cumulative impacts would be expected from the proposed activities or as the result of past, present or reasonably foreseeable oil and gas leasing, exploration, development and production in the GOM.

### **Conclusion**

The sections above discuss the potential range of effects to sea turtles from the proposed action, including: (1) vessel noise and collisions; (2) marine debris; (3) water-quality degradation from drilling rig effluents; (4) oil spills and spill-response activities; and (5) drilling noise. The potential effects of the proposed activity on sea turtles will not rise to the level of significance for the following reasons:

- The best available scientific information indicates that sea turtles do not greatly use sound in the environment for survival; therefore, disruptions in environmental sound would have little effect.
- The scope, timing, and transitory nature of the proposed action will produce limited amounts of drilling noise in the environment. As described, effects of vessel noise on sea turtles are considered “discountable” (USDOC, NMFS, 2020a).
- Implementation of Appendix B: Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols and the International Convention for the Prevention of Pollution from Ships (MARPOL) requirements under the United States Coast Guard (USCG), appreciably reduces the likelihood of sea turtles encountering marine debris from the proposed activity.

The risk of collisions between sea turtles and vessels associated with the proposed action exists but would not rise to the level of significance given:

- Under 30 CFR § 550.282, as clarified by mitigation and monitoring measures in Appendix C: Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols outlined in the NMFS 2020 BO , BOEM provides guidelines for the monitoring programs designed to minimize the risk of vessel strikes to sea turtles and other protected species and the reporting of any observations of injured or dead protected species.
- The NMFS 2020 BO determined that monitoring measures should appreciably reduce the potential for vessel strikes. The NMFS issued an Incidental Take Statement on sea turtle species; the Statement contains reasonable and prudent measures (RPMs) with implementing terms and conditions to help minimize take. As the operator has indicated that the vessel strike avoidance guidance (Appendix C: Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols) will be followed, there should be appreciably reduced numbers of sea turtles that may be incidentally taken from routine offshore vessel operations; however, the available information on the relationship between these species and OCS oil and gas activities indicates that sea turtles may be killed or injured by vessel strikes. Therefore, pursuant to Section 7(b)(4) of the ESA, NMFS anticipates incidental take and granted a limited number of Incidental Take Authorizations to BOEM for sea turtle mortalities by vessel strikes. BOEM continues to monitor for any strikes to ensure this authority is not exceeded and to date, none have been reported. The scope, timing, and transitory nature of the proposed action will result in limited opportunity for vessel strikes to sea turtles during operations.

### **3.6.2.3. Alternative 3**

If selected, Alternative 3, the Proposed Action with Additional Conditions of Approval, would allow the operator undertaking the proposed activities, as requested and conditioned in the plan; however, the operator would be required to undertake additional mitigation and monitoring measures as identified by BOEM and NMFS. Alternative 3 differs from Alternative 2 because the additional mitigation and monitoring measures address this resource. NMFS provided concurrence, in that, the proposed activities and the use of a moon pool are within the scope of the effect analysis of the NMFS 2020 BO. Because the operator is required to follow all existing lease stipulations and regulations as clarified by NTLs, conditions outlined in the previous analyses related to BOEM NTL Nos. 2015-G03, 2016-G01, and 2016-G02; and the requirement for additional mitigation and monitoring measures, should minimize or reduce the risk of a significant impact to sea turtles.

## **3.7. FISH RESOURCES AND ESSENTIAL FISH HABITAT**

### **3.7.1. Affected Environment**

A detailed description of the Fish Resources and Essential Fish Habitat (EFH) of the GOM may be found in Chapter 4.7 of the Multisale EIS and 2018 SEIS, and is incorporated by reference into this SEA. The following section provides a summary of the information found in the Multisale EIS and 2018 SEIS.

The NMFS 2020 BO identified the following Federally listed threatened fish species in the GOM: the Gulf sturgeon, the oceanic whitetip shark and the giant manta ray. The Gulf sturgeon (*Acipenser oxyrinchus oxyrinchus*) was listed as threatened October 30, 1991 (56 CFR §49653, September 30, 1991). The oceanic whitetip shark (*Carcharhinus longimanus*) was listed as threatened January 30, 2018 under the ESA (83 FR 4153). The giant manta ray (*Manta birostris*) was listed as threatened January 22, 2018 under the ESA

(83 FR 2916). A detailed description of the Gulf sturgeon and critical habitat, and oceanic white tip shark and giant manta ray may be found in Sections 6.2.11 to 6.2.14 of the NMFS 2020 BO.

The NMFS and USFWS jointly designated Gulf sturgeon critical habitat on April 18, 2003 (50 CFR §226.214). Seven of the marine and estuarine units of the Gulf sturgeon critical habitat (Units 8-14) are found along the states of Louisiana, Mississippi, Alabama, and Florida (Figure 90 of the NMFS 2020 BO).

EFH is defined as “those waters and substrate necessary to fish for spawning, breeding, feeding, and growth to maturity” [16 U.S.C. § 1801(10)]. These habitats are crucial for maintaining healthy fish resources and fishery stocks. Due to the wide variation of habitat requirements for all life history stages of managed species, NOAA and the Gulf of Mexico Fishery Management Council initially identified EFH throughout the GOM to include all coastal and marine waters and substrates from the shoreline to the seaward limit of the Exclusive Economic Zone (200 mi [322 km] from shore). The EFH final rule summarizing EFH regulation (50 CFR § 600) outlines additional interpretation of the EFH definition. Waters, as defined previously, include “aquatic areas and their associated physical, chemical, and biological properties that are used by fish, and may include aquatic areas historically used by fish where appropriate.” Substrate includes “sediment, hard bottom, structures underlying the waters, and associated biological communities.” Necessary is defined as “the habitat required to supporting a sustainable fishery and the managed species contribution to a healthy ecosystem.” “Fish” includes “finfish, mollusks, crustaceans, and all other forms of marine animal and plant life other than marine mammals and birds,” whereas “spawning, breeding, feeding or growth to maturity” covers the complete life cycle of those species of interest.

The EFH for the oceanic whitetip shark in the project area includes localized areas in the central GOM and Florida Keys. Although no EFH or critical habitat has been designated, the giant manta rays are widespread in the GOM. Giant manta rays occupy tropical, subtropical, and temperate oceanic waters and productive coastlines and are commonly found offshore in oceanic waters, but are sometimes found feeding in shallow waters (less than 10 meters) during the day (Miller, 2016).

The GOM supports a great diversity of fish species, including a wide variety of commercially and recreationally valuable fishes, most of which are linked either directly or indirectly to the estuaries ringing the Gulf. The life history of estuarine-dependent species involves spawning on the continental shelf; the transportation of eggs, larvae, or juveniles back to the estuary nursery grounds; and the migration of the adults back to the sea for spawning. Monthly ichthyoplankton collections over the years 2004-2006 offshore of Alabama confirmed that peak seasons for ichthyoplankton concentrations on the shelf are spring and summer (Hernandez et al., 2010). Additionally, the waters of the northern GOM support many coastal pelagic fishes and highly migratory species, some of which spawn exclusively in this region. The distribution of fish species is related to ecological factors (e.g., salinity, temperature, bottom type, primary production and availability of prey) which vary, sometimes widely, across the Gulf and between inshore and offshore waters. Characteristic fish resources are associated with various environments and are not randomly distributed.

Although a generalized analysis suggests, for locations off the continental shelf, species richness and abundance decrease with depth, Rowe and Kennicutt (2009) found food resources are a dominant factor controlling distribution of deepwater benthos in the GOM. Inputs such as the Mississippi River and hydrocarbon seep communities influence local densities of fauna associated with a given depth zone. Descriptions of ecological groups of fishes that occur in the region, including oceanic pelagics and mesopelagics, can be found in Chapter 4.7 of the Multisale EIS and 2018 SEIS.

### **3.7.2. Impact Analysis**

A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed exploration activities on fish and essential fish habitat can be found in Chapters 4.7 of the Multisale EIS and 2018 SEIS, and is incorporated by reference. A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed exploration activities on Federally listed threatened species can be found in Sections 8.4, 8.7, 8.8, 10, 11.3 and 11.4 of the NMFS 2020 BO, and is incorporated by reference. The IPFs associated with the proposed activities that could affect EFH and fish resources include: (1) coastal and marine environmental degradation; (2) presence of a MODU; (3) temporary discharge of drilling cuttings and associated drilling fluids; and (4) blowouts and oil spills. The NMFS 2020 BO identified IPFs associated with activities proposed that could affect federally listed threatened species including: (1) marine trash and debris; (2) vessel collisions; and (3) oil spills.

### **3.7.2.1. Alternative 1**

If selected, Alternative 1, No Action Alternative, would result in the operator not undertaking the proposed activities as described in the plan. Therefore, the site-specific IPFs to fish and EFH would not occur. For example, there would be no drilling noise that would result in behavioral change, masking, or non-auditory effects to the fish resources, no long-term or permanent displacement of fish resources from preferred habitats, and no destruction or adverse modification of any habitats. Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas exploration and development activities as described in the Multisale EIS and 2018 SEIS and routine, accidental, and cumulative impacts would continue to occur.

### **3.7.2.2. Alternative 2**

If selected, Alternative 2, the Proposed Action as Submitted and in accordance with their lease stipulations (in addition to any amendments), OCSLA requirements, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), Reasonable and Prudent Measures and Implementing Terms and Conditions as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and with FWS that concluded April 20, 2018 (FWS 2018 BO). However, no additional, site-specific mitigation and monitoring measures would be required by BOEM or NMFS. As described in the analyses below, impacts to fish and EFH from the proposed action are expected to be short-term, localized and not lead to significant impacts. Impacts from the proposed activities are not expected to reduce appreciably, the likelihood of both the survival and recovery of Federally listed threatened species.

### **Routine Activities**

Routine activities, such as the discharge of drilling fluids and cuttings offshore would contribute to localized temporary marine environmental degradation. Drilling operations are restricted in time, and pelagic species in the area could easily avoid discharge plumes. Routine discharges from the MODU would be highly diluted in the open marine environment. The presence of the MODU will act as a fish-attracting device for the short period of time the rig is on site; however, routine discharges on fish resources will be very limited in duration.

In the last five-year NMFS species review, vessel strikes were identified as an emerging threat for Gulf sturgeon. The NMFS 2020 BO *Effects Analysis* for Gulf sturgeon estimated one nonlethal and 21 lethal vessel strikes would occur over 50 years as a result of vessels associated with the proposed action. The effects to giant manta rays and oceanic white tip sharks from vessel strikes are discountable. The operator is required to adhere with the mitigation and monitoring measures provided in the NMFS 2020 BO under Appendix C: Gulf of Mexico Vessel Strike Avoidance and Injured/Dead Aquatic Protected Species Reporting Protocols. Compliance with the regulations as clarified in the NMFS 2020 BO should reduce or avoid impacts from vessel strikes under this alternative. Many types of plastic materials end up as solid waste during drilling and production operations. Some of this material is accidentally lost overboard where fish or federally listed threatened species could consume it or become entangled in it. The incidental ingestion of marine debris and entanglement could adversely affect fish and Federally listed threatened species. Implementation of Appendix B: Gulf of Mexico Marine Trash and Debris Awareness and Elimination Survey Protocols and MARPOL requirements under the USCG may reduce, but not eliminate the risk of marine debris (USDOC, NMFS, 2020a). The NMFS 2020 BO determined that Gulf sturgeon, oceanic whitetip sharks, and giant manta rays are not likely to be adversely affected by marine debris resulting from the proposed action.

### **Accidental Events**

Accidental blowouts and spills with limited quantities of hydrocarbons also have the potential to affect fish resources and EFH, but there is no evidence to date that fish or EFH in the Gulf have been adversely affected at a population level by spills or chronic contamination. At the scale of this EA, any accidental impact would be limited in scope and affected fishes would likely be replaced by organisms from beyond the area of impact or would be colonized during the next recruitment event. Early life stages of fishes may be more sensitive than adults to potentially adverse impacts resulting from exposure to hydrocarbons. For

this reason, BOEM considers eggs and larval fishes to be at greater risk than adults in the event of exposure to contamination resulting from a spill or blowout. The specific effects of oil on fish can include direct lethal toxicity, sublethal disruption of physiological processes (internal lesions), suffocation due to oil coating gills, incorporation of hydrocarbons causing tainting or accumulation in the food chain, and changes in biological habitat (Moore and Dwyer, 1974; Incardona et al., 2014; Murawski et al., 2014). However, due to typically high fecundity and relatively wide distribution of eggs and larvae, it is unlikely spilled contaminants would overlap spatially and temporally with a fraction of eggs and larvae large enough to significantly impact populations. Furthermore, most adult fishes are expected to avoid adverse environmental conditions, minimizing the potential for impacts resulting from oil and dispersants. Estuaries are important nursery areas (EFH) for fish and aquatic life. Impacts related to oiling of these areas could result in the destruction of marsh habitat, facilitate in the erosion of coastlines, and increase the potential for adversely impacting juvenile fishes. A discussion of the impacts of oil on adult fish, fish eggs, and larvae can be found in Chapter 4.7 of the Multisale EIS and 2018 SEIS. Given that the potential for a blowout or a small spill, there is a limited possibility for large amounts of oil released from a blowout or spill reaching shore. Additional sensitive habitat features and potential impacts to these habitats are discussed in **Chapter 3.4** (Deepwater Benthic Communities) of this document.

Accidental blowouts and spills also have the potential to affect federally listed species, but there is no evidence to date that there have been adverse impacts to population levels by spills or chronic contamination. Considering the location of Gulf sturgeon critical habitat in relation to oil and gas activities, the likely dilution of oil reaching nearshore areas, and the on-going weathering and dispersal of oil over time, it is not anticipated that the effects from oil spills will appreciably diminish the value of Gulf sturgeon designated critical habitat for the conservation of the species.

The likelihood of an individual giant manta ray or oceanic white tip shark being in the area of an oil spill is small and only those individuals found in the footprint of an oil spill would be affected. A small number of giant manta rays or oceanic white tip sharks are likely to be exposed to oil, and those exposures would likely result in effects similar to other marine species. Because data related to abundance estimates for oceanic whitetip sharks and giant manta rays in the GOM is limited, NMFS is not able to quantify an estimated number of oil spill exposures or mortalities for this species.

## **Cumulative Impacts**

Cumulative activities that could impact fish, EFH, and Federally listed threatened species in the area of the proposed action include State oil and gas activity, coastal development, crude oil imports by tanker, commercial and recreational fishing, hypoxia (i.e., red or brown tides), removal of OCS structures, vessel strikes, and offshore discharges of drilling muds and produced waters. It is expected that environmental degradation from the proposed action and non-OCS activities would affect fish populations, EFH, and Federally listed threatened species; however, the incremental contribution of the proposed action to these cumulative impacts would be small and almost undetectable for fish and EFH and discountable for Federally listed threatened species. Therefore, no significant cumulative impacts on EFH, fish resources, or Federally listed threatened species would be expected as a result of the proposed activities when added to the impacts of past, present, or reasonably foreseeable oil and gas development in the area as well as other activities in the area.

## **Conclusion**

The proposed action is expected to have little impact on any fish, EFH, or Federally listed threatened species endemic to the northern GOM. Specific effects from any one oil spill would depend on several factors, including timing, location, volume and type of oil, environmental conditions, and countermeasures used. If a blowout occurred, ichthyoplankton, fish eggs, or larvae would suffer mortality in areas where their numbers are concentrated and where oil concentrations are high. However, impacts are still expected to be minimal to nonexistent based on the low probability of a spill occurring (see **Chapter 1.4**).

### **3.7.2.3. Alternative 3**

If selected, Alternative 3, the Proposed Action with Additional Conditions of Approval, would result in the operator undertaking the proposed activities, as requested and conditioned in the plan; however, the operator would be required to undertake additional condition(s) of approval as identified by BOEM. As described in the analyses above, impacts to fish, EFH, and Federally listed threatened species from the proposed action are expected to be short-term, localized and not lead to significant impacts. The condition

of approval outlined in **Chapter 2.4** is not expected to increase or decrease the potential for effects to fish, EFH, or Federally listed threatened species from the proposed action. Alternative 3 does not differ from Alternative 2 because the additional mitigation and monitoring measures do not address this resource (i.e., all assumptions, estimates, and conclusions are identical); see the analysis provided in **Chapter 3.7** for Alternative 2 for this resource.

## **3.8. ARCHAEOLOGICAL RESOURCES**

### **3.8.1. Affected Environment**

Archaeological resources are defined in 30 CFR § 550.105 as “any material remains of human life or activity that are at least 50 years of age and that are of archaeological interest.” Archaeological resources on the OCS can be divided into two types: prehistoric and historic. Detailed descriptions of these resource types are provided in Chapter 4.13 of the Multisale EIS and 2018 SEIS. The following information is a summary of these descriptions, which are incorporated by reference into this SEA.

#### *Prehistoric*

Geologic features that have a high probability for associated prehistoric sites in the northwestern and north central Gulf (from Texas to Alabama) include barrier islands, river channels and associated floodplains and terraces, and salt dome features. Also, a high probability for prehistoric resources may exist landward of a line that roughly follows the 60 m (197 ft) bathymetric contour, which represents the Pleistocene shoreline during the last glaciation some 12,000 years ago when the coastal area of Texas and Louisiana is generally considered to have been populated. BOEM is reviewing evidence to determine if a change in the currently accepted area of prehistoric site probability is warranted. The water depth in the area of the proposed action precludes the potential for prehistoric sites or artifacts.

#### *Historic*

Historic archaeological resources on the federal OCS include shipwrecks and a single light house (Ship Shoal Light). Historic research has identified over 4,000 potential shipwreck locations in the Gulf, with nearly 1,500 of these potential shipwreck locations on the OCS (Garrison et al., 1989). The historic record, however, is by no means complete, and the current ability to predict potential sites has proven inaccurate. As demonstrated by several studies (Pearson et al., 2003; Lugo-Fernandez et al., 2007; Krivor et al., 2011; Rawls and Bowker-Lee, 2011), many more shipwrecks are likely to exist on the seafloor than have been accounted in available historic literature. Currently a high-resolution remote sensing survey is the most reliable method for identifying and avoiding historic archaeological resources.

A 2003 study recommended including some deepwater areas, primarily on the approach to the Mississippi River, among those lease areas requiring archaeological investigation. With this in mind, BOEM revised its guidelines for conducting archaeological surveys in 2005 and added about 1,200 lease blocks to the list of blocks requiring an archaeological survey and assessment. Archaeological survey blocks were further expanded in 2011 and current requirements are posted on the BOEM website under NTL No. 2005-G07 and Joint NTL No. 2011-G01. At present, high-resolution geophysical, ROV, and/or diver survey and investigation is required for bottom disturbing activities.

Historic shipwrecks have, with the exception of three significant vessels found by treasure salvors, been primarily discovered through oil industry sonar surveys in water depths up to 9,000 ft (2,743 m). In the last five years, over four dozen potential shipwrecks have been located and several of these ships have been confirmed visually as historic vessels. Many of these wrecks were not previously suspected to exist in these areas, based on the historic record. The preservation of historic wrecks found in deep water has been outstanding because of a combination of environmental conditions and limited human access.

The *Deepwater Horizon* spill released an estimated 53,000-62,000 bbl of oil per day for almost three months. Much of the oil was treated with dispersant at the sea surface and at the source in a water depth of 5,000 ft (1,524 m). The use of dispersants at the wellhead could result in currently unknown effects from dispersed oil droplets settling to the seafloor and could possibly contaminate exposed artifacts and wood or steel hulls such as those observed on many deepwater sites (Atauz et al., 2006; Church et al., 2007; Church and Warren, 2008; Ford et al., 2008).

The best available information does not provide a complete understanding of the effects, if any, of the spilled oil and potential response/cleanup activities on archaeological resources that may be located in deep

water. Though information on the actual impacts to submerged archaeological resources is non-existent at this time, oil settling to the seafloor due to dispersant use at the wellhead could come into contact with archaeological resources. At present, there is no evidence of this having occurred. An experimental study has suggested that while the degradation of wood in terrestrial environments is initially retarded by contamination with crude oil; at later stages, the biodeterioration of wood is accelerated (Ejechi, 2003). While there are different environmental constraints that affect the degradation of wood in terrestrial and waterlogged environments, soft-rot fungal activity, one of the primary wood degrading organisms in submerged environments, was shown to be increased in the presence of crude oil.

### **3.8.2. Impact Analysis**

A detailed impact analysis of the routine, accidental, and cumulative impacts of the proposed activities on historic archeological resources can be found in Chapters 4.13 the Multisale EIS and 2018 SEIS, and is incorporated by reference. The IPF associated with the proposed action that could affect submerged archaeological resources is seafloor disturbance. These discussions also are summarized below and incorporated by reference into this SEA.

The routine IPF associated with Anadarko's proposed development activities in the area of the proposed action that could affect archaeological resources is limited to direct contact or disturbance during well emplacement activities or equipment used for the drilling operations.

The historically available literature is not sufficient to identify historic shipwreck losses in the area of the proposed action as historic records of losses occurring this far offshore are not location-specific (Pearson et al., 2003; Lugo-Fernandez et al., 2007; Krivor et al., 2011; and Rawls and Bowker-Lee, 2011). However, if a historic resource exists in the area of drilling, direct physical contact with a shipwreck site could destroy fragile materials, such as the hull remains or artifacts, and could disturb the site context (Atauz et al., 2006; Church and Warren, 2008). To date, two historically significant shipwrecks were found to have suffered damage from drilling activities because of a lack of knowledge of their presence.

The IPFs that could be associated with accidental events include seafloor disturbances from jettisoned/lost debris and, as discussed above, deterioration from potential oil spills. Similar to routine impacts, discarded/lost material that falls to the seabed has the potential to damage and/or disturb archaeological resources. Oil spills and their remediation efforts could also accelerate deterioration of archaeological resources. A detailed discussion of all potential impacts is found below.

#### **3.8.2.1. Alternative 1**

If selected, Alternative 1, the No Action Alternative, would result in the operator not undertaking the proposed activities as described in the plan. Therefore, the IPFs mentioned above (i.e., bottom disturbance associated with well emplacement and the use of equipment associated with drilling operations) would not take place, and any impact that these actions could cause would not occur. Likewise, under the No Action Alternative, there would be no possibility of a spill. As a result, whatever archaeological resources may be present in the area of potential effect (APE) would not be affected in any way if the No Action Alternative was selected. Activities related to previously issued leases and permits (as well as those that may be issued in the future under a separate decision) related to the OCS activities would not increase. The No Action Alternative would not significantly change the environmental impacts of overall OCS oil and gas exploration and development activities as described in the Multisale EIS and 2018 SEIS and routine, accidental, and cumulative impacts would continue to occur.

#### **3.8.2.2. Alternative 2**

If selected, Alternative 2, the Proposed Action as Submitted and in accordance with their lease stipulations (in addition to any amendments), OCSLA requirements, and all applicable regulations (as per 30 CFR § 550.101(a)), and guidance provided in all appropriate NTLs (as per 30 CFR § 550.103), Reasonable and Prudent Measures and implementing Terms and Conditions as required under the ESA Consultation with NMFS that concluded March 13, 2020 (NMFS 2020 BO) and with FWS that concluded April 20, 2018 (FWS 2018 BO). However, no additional, site-specific mitigation and monitoring measures would be required by BOEM or NMFS. Examples of potential impacts to archaeological resources would include, but are not limited to, damage to potential resources from well emplacement activities, lost/discarded material, and potential impacts from an accidental oil spill. As described in the proposed plan and discussed below, the proposed activities are not expected to have significant impacts on known or unknown historical archaeological resources.

## Routine Activities

Impacts to a historic site could result from direct physical contact causing irreversible damage. The undisturbed provenience of archaeological data (i.e., the 3D location of archaeological artifacts) allows archaeologists to accumulate a record of where every item is found, and to develop a snapshot as to how artifacts relate to other items or the site as a whole. The analysis of artifacts and their provenience is one critical element used to make a determination of eligibility to the National Register of Historic Places and is essential in understanding past human behavior and ways of life. Impacts from the proposed operations could alter the provenience and destroy fragile remains, such as the hull, wood, glass, ceramic artifacts and possibly even human remains, or information related to the operation or purpose of the vessel. The destruction and loss of this data eliminate the ability of the archaeologist to fully and accurately detail activity areas found at the site, variation and technological advances lost to history, the age, function, and cultural affiliation of the vessel, and its overall contribution to understanding and documenting the maritime heritage and culture of the region.

BOEM's regulation at 30 CFR § 550.194 requires that an archaeological survey be conducted prior to development of leases if the Regional Director has reason to believe that an archaeological resource may exist in the lease area. An Archaeological survey has been conducted in Mississippi Canyon Blocks 80, 81, 82, 125, and 126. No targets that may represent significant archaeological resources were identified in the high-resolution geophysical survey near/within the area of Anadarko's proposed activity.

## Accidental Events

Although unlikely, accidental blowouts and spills from the proposed action could lead to oil contact with submerged archaeological resources. While there is no information on the actual impacts of the *Deepwater Horizon* spill on submerged archaeological resources, should an accidental blowout and spill occur during the operator's proposed action, oil may settle on the seafloor due to dispersant use at the wellhead and could come into contact with archaeological resources. Although there is uncertainty and limited data on the effects of an oil spill at depth on submerged archaeological resources, an experimental study has suggested that while the degradation of wood in terrestrial environments is initially retarded by contamination with crude oil; at later stages, the biodeterioration of wood is accelerated (Ejechi, 2003). While there are different environmental constraints that affect the degradation of wood in terrestrial and waterlogged environments, soft-rot fungal activity, one of the primary wood degrading organisms in submerged environments, was shown to be increased in the presence of crude oil. No impacts are expected from marine remediation efforts because bottom-disturbing activities are not anticipated due to the water depth.

Another IPF that could result from an accidental event is from the loss of debris from the MODU during drilling operations. Debris such as structural components (i.e., grating, wire, tubing, etc.), boxes, pallets, and other loose items can become dislodged during heavy seas or storm events and fall to the seabed. Similarly, thousands of joints of drill pipe are used during drilling operations; requiring regular transport out to the MODU via workboats. There is the potential to lose pieces of drill pipe during transfer operations or when "tripping pipe" in and out of the wellbore. Similar to the impacts noted under Routine Activities, if lost drill pipe or debris were to fall onto an unknown archaeological resource near the well site, damage could destroy fragile materials, such as the hull remains and artifacts, and could disturb the site's context and associated artifact assemblage. Additionally, lost material could result in the masking of actual archaeological resources or the introduction of false targets that could be mistaken in the remote sensing record as historic resources.

In the event of a catastrophic spill similar to the *Deepwater Horizon* spill, any substantive impact to archaeological resources is very unlikely because the potential impacts from a catastrophic spill would be similar to aforementioned routine and accidental issues. However, despite the recent *Deepwater Horizon* spill, historical trends in the GOM (see **Chapter 1.4**) indicate that catastrophic spill events are not likely to occur as a result of the proposed action.

## Cumulative Analysis

Cumulative impacts on unknown archaeological resources that may be present in the area of the proposed action could result from other GOM activities. Since the water depth at the proposed well sites ranges from approximately 3,710 to 4,350 ft (1,131 to 1,326 m) and the area of the proposed action is over

53 mi (85 km) from shore, those activities would be limited to commercial fishing, marine transportation, and adjacent oil and gas exploration, development, and production operations.

During adjacent oil and gas operations, commercial fishing, and maritime transportation activities, there is associated the loss or discard of debris that could result in the masking of archaeological resources or the introduction of false targets that could be mistaken in the remote sensing record as historic resources. Future exploration, development, and production operations and/or any related infrastructure support could lead to bottom disturbances in the area of the proposed action; however, no additional activities have been proposed or are under review at this time. Similarly, G&G surveys have been permitted near the area of the proposed action. These surveys may involve the seabed deployment of receivers attached to degradable concrete anchors that are deployed from the sea surface. These anchors have the potential to damage unknown archaeological resources that may exist in the area of the proposed action as they descend through the water column; however, their small size and relatively light weight (~65 lbs [34 kg]) is not expected to cause significant impacts.

Any known or unknown archaeological resources that may be present in Mississippi Canyon Blocks 80, 81, 82, 125, and 126 could be impacted by contact with oil from a blowout or spill from adjacent oil and gas operations. Similarly, cumulative impacts from accidental oil spills and remediation efforts for adjacent oil and gas operations are not expected because of the water depth at the proposed well sites and the historically low probability of a loss of well control/blowout.

Considering the potential cumulative impacts from all other GOM activities, the operator's proposed activities would constitute the primary effect, if any, on any known or unknown archaeological resource that may exist in the area of the proposed action. However, based on the archaeological assessment conclusions, there is no reason to believe that the proposed action would result in the disturbance of archaeological resources. Therefore, no significant cumulative impacts are expected as a result of the proposed action when added to the impacts of past, present, or reasonably foreseeable oil and gas development in the area as well as other proximal activities.

### **3.8.2.3. Alternative 3**

If selected, Alternative 3, the Proposed Action with Additional Conditions of Approval, would result in the operator undertaking the proposed activities, as requested and conditioned in the plan; however, the operator would be required to undertake additional mitigation and monitoring measures as identified by BOEM and NMFS. As described in the analysis above, impacts to archaeological resources are not expected. The condition of approval outlined in **Chapter 2.4** is not expected to increase or decrease the potential for effects to archaeological from the proposed action. Alternative 3 does not differ from Alternative 2 because the additional mitigation and monitoring measures do not address this resource (i.e., all assumptions, estimates, and conclusions are identical).

### **Conclusion**

Based on the previous information and the survey conclusions, there is no reason to believe that archaeological resources could be present in the area of the proposed action. If an unknown archaeological resource were to exist where bottom-disturbing operations are proposed to occur, and the operator were unaware of its existence prior to disturbing the bottom, the operator's activities might have a significant impact on that resource. Such impact would be damage and/or disturbance to the resource from drilling the well and from the associated equipment. Impacts from accidental events related to the proposed action such as accidental oil spills and their remediation efforts are not expected because of the water depth at the well sites and the historically low probability of a loss of well control/blowout. However, debris resulting from accidental events could lead to impacts similar to those expected from routine impacts such as contact with the well and/or well equipment.

## **4. CONSULTATION AND COORDINATION**

The Endangered Species Act of 1973 (ESA) (16 U.S.C. §§ 1531 *et seq.*), as amended, establishes a national policy designed to protect and conserve threatened and endangered species and the ecosystems upon which they depend. Section 7(a)(2) of the ESA requires each Federal agency to ensure that any action that they authorize, fund, or carry out is not likely to jeopardize the continued existence of a listed species or result in the adverse modification of designated critical habitat. On April 20, 2018, the U.S. Fish and Wildlife Service (FWS) issued its 10-year programmatic BO for BOEM and BSEE's oil and gas activities

in the GOM. The FWS 2018 BO does not include any terms and conditions for the protection of endangered species that the Bureaus, lessees, or operators must implement but did provide recommendations for Oil Spill Contingency Plan information, aircraft, marine debris and trash, information needs, and future coordination. The FWS 2018 BO also noted that any future consultations may be informal, dependent upon the likelihood of take.

On March 13, 2020, NMFS issued a BO and related terms and conditions for oil and gas activities in the GOM for the protection of these species, including holding lease sales. The NMFS programmatic BO addresses any future lease sales and any approvals issued by BOEM and BSEE, under both existing and future OCS oil and gas leases in the GOM, over a 10-year period. Applicable terms and conditions and reasonable and prudent measures from the NMFS 2020 BO will be applied at the lease sale stage; other specific conditions of approval will also be applied to post-lease approvals. The NMFS 2020 BO may be found here: <https://www.fisheries.noaa.gov/resource/document/biological-opinion-federally-regulated-oil-and-gas-program-activities-gulf-mexico>. The Appendices and protocols may be found here: <https://www.fisheries.noaa.gov/resource/document/appendices-biological-opinion-federally-regulated-oil-and-gas-program-gulf-mexico>. BOEM petitioned NMFS for rulemaking under the MMPA, to assist industry in obtaining incidental take coverage for marine mammals due to oil and gas and G&G surveys in the GOM. If NMFS issues a final rule as a result of the petition, the NMFS programmatic BO may be amended and additional mitigation measures beyond what is currently within Appendix A and C may be imposed through Letters of Authorizations under the rulemaking for MMPA.

In accordance with the National Historic Preservation Act (54 U.S.C. §§ 300101 *et seq.*), Federal agencies are required to consider the effects of their undertakings on historic properties. The implementing regulations for Section 106 of the National Historic Preservation Act, issued by the Advisory Council on Historic Preservation (36 CFR § 800), specify the required review process. In accordance with 36 CFR § 800.8(c), BOEM intends to use the NEPA substitution process and documentation for preparing an EIS/ROD or an EA/FONSI to comply with Section 106 of the National Historic Preservation Act in lieu of 36 CFR §§ 800.3-800.6

## 5. PUBLIC COMMENT

Once the operator's plan was deemed submitted (as per 30 CFR § 550.231) on September 10, 2020, and it was placed on <http://www.regulations.gov> for a 10-day public review. At the end of the comment period on September 21, 2020, no comments were received.

## 6. REFERENCES

- American Petroleum Institute (API). 1989. Effects of offshore petroleum operations on cold water marine mammals: A literature review. Washington, DC: American Petroleum Institute. 385 pp.
- Anadarko Petroleum Corporation (Anadarko). 2020. Initial Exploration Plan, N-10117. Mississippi Canyon Block 80, Lease Number OCS-G35311, Mississippi Canyon Block 81, Lease Number OCS-G35312, Mississippi Canyon 82, Lease Number OCS-G35313, Mississippi Canyon Block 125, Lease Number OCS-G35316, and Mississippi Canyon Block 126, Lease Number OCS-G18194. Offshore Alabama.
- Atauz, A.D., W. Bryant, T. Jones, and B. Phaneuf. 2006. Mica shipwreck project: Deepwater archaeological investigation of a 19th century shipwreck in the Gulf of Mexico. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2006-072. 116 pp.
- Australian Maritime Safety Authority (AMSA). 2003. Post Spill Monitoring: Background Paper. Prepared by Wardrop Consulting and the Cawthron Institute for the Australian Maritime Safety Authority (AMSA) and the Marine Safety Authority of New Zealand (MSA). Published by AMSA, Canberra. ISBN: 0 642 709912.
- Basan, P.B., C.K. Chamberlain, R.W. Frey, J.D. Howard, A. Seilacher and J.E. Warne. 1978. Trace fossil concepts. Society of Economic Paleontologists and Mineralogists, Short Course No. 5, Tulsa, OK. 181 pp.

- Bright, T.J. and R. Rezak. 1978. Northwestern Gulf of Mexico topographic features study: Final report. U.S. Dept. of the Interior, Bureau of Land Management, New Orleans OCS Office, New Orleans, LA. Study No. 1978-4. 692 pp. Internet website: <https://espis.boem.gov/final%20reports/4069.pdf> Accessed September 23, 2020.
- Bright, T.J., R. Rezak, A.H. Bouma, W.R. Bryant, and W.E. Pequegnat. 1976. A biological and geological reconnaissance of selected topographical features on the Texas continental shelf. Final report to the U.S. Dept. of the Interior, Bureau of Land Management, New Orleans OCS Office, New Orleans, LA. Contract # 08550-CT5-4.
- Church, R.A. and D.J. Warren. 2008. Viosca Knoll wreck: Discovery and investigation of an early nineteenth-century sailing ship in 2,000 feet of water. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2008-018. 41 pp.
- Church, R., D. Warren, R. Cullimore, L. Johnston, W. Schroeder, W. Patterson, T. Shirley, M. Kilgour, N. Morris, and J. Moore. 2007. Archaeological and biological analysis of World War II shipwrecks in the Gulf of Mexico: Artificial reef effect in deep water. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2007-015. 387 pp.
- Coastal Response Research Center. 2007. Submerged oil – State of the practice and research needs. Prepared by the Coastal Response Research Center, Durham, NH. 29 pp. + app. Conn, P. B., and G. K. Silber. 2013. Vessel speed restrictions reduce risk of collision-related mortality for North Atlantic right whales. *Ecosphere* 4(4):1–16.
- Conn, P. B., and G.K. Silber. 2013. Vessel speed restrictions reduce risk of collision-related mortality for North Atlantic right whales. *Ecosphere* 4(4):1–16.
- Continental Shelf Associates, Inc. (CSA). 2006. Effects of oil and gas exploration and development at selected continental slope sites in the Gulf of Mexico. Volume II: Technical report. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2006-045. 636 pp.
- Cordes, E.E., M.P. McGinley, E.L. Podowski, E.L. Becker, S., Lessard-Pilon, S.T. Viada, and C.R. Fisher. 2008. Coral communities of the deep Gulf of Mexico. *Deep-Sea Research* 55(6):777-787.
- Davis, R.W., W.E. Evans, and B. Würsig. 2000. Cetaceans, sea turtles and seabirds in the northern Gulf of Mexico: Distribution, abundance and habitat associations. Volume II: Technical report. U.S. Dept. of the Interior, Geological Survey, Biological Resources Division, USGS/BRD/CR-1999-0006 and U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2000-003. 346 pp.
- Eckdale, A.A., R.G. Bromley, and S.G. Pemberton. 1984. Ichnology: The use of trace fossils in sedimentology and stratigraphy. Society of Economic Paleontologists and Mineralogists, Short Course No. 15, Tulsa, OK. 317 pp.
- Ejechi, B.O. 2003. Biodegradation of wood in crude oil polluted soil. *World Jour. Microb. & Biotech.* 19:799–804.
- Federal Register*. 2014. Endangered and threatened species: Critical habitat for the northwest Atlantic Ocean loggerhead sea turtle distinct population segment (DPS) and determination regarding critical habitat for the North Pacific Ocean loggerhead DPS. Final rule (50 CFR 256). 79 FR 39855, pp. 39855-39912.
- Federal Register*. 2016a. Endangered and Threatened Wildlife and Plants: 12-Month Finding on a Petition to Downlist the West Indian Manatee, and Proposed Rule to Reclassify the West Indian Manatee as Threatened; Proposed Rule. Proposed Rules (50 CFR § 17). 81 FR 999, pp. 1000-1026.
- Federal Register*. 2016b. Endangered and Threatened Wildlife and Plants: Notice of 12-Month Finding on a Petition to List the Gulf of Mexico Bryde's Whale as Endangered Under the Endangered Species Act (ESA); Proposed Rule. 81 FR 88639, pp. 88639-88656.

- Federal Register*. 2017. Endangered and Threatened Wildlife and Plants: Reclassification of the West Indian manatee From Endangered to Threatened (50 CFR § 17). 82 FR 16668, pp. 16668-16704.
- Federal Register*. 2019. Endangered and Threatened Wildlife and Plants: Endangered Status of the Gulf of Mexico Bryde's Whale (50 CFR § 224). 84 FR 15446, pp. 15446-18488.
- Fisher, C.R. 1990. Chemoautotrophic and methanotrophic symbioses in marine invertebrates. *Reviews in Aquatic Sciences* 2:399-436.
- Fisher, C., H. Roberts, E. Cordes, and B. Bernard. 2007. Cold seeps and associated communities of the Gulf of Mexico. *Oceanography* 20(4):118-129.
- Ford, B., A. Borgens, W. Bryant, D. Marshall, P. Hitchcock, C. Arias, and D. Hamilton. 2008. Archaeological excavation of the Mardi Gras Shipwreck (16GM01), Gulf of Mexico continental slope. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Report MMS 2008-037. 313 pp.
- French-McCay, D.P. 2004. Oil spill impact modeling: development and validation. *Environmental Toxicology and Chemistry*, 23(10), 2441-2456.
- Frey, R.W. 1975. The study of trace fossils: A synthesis of principles, problems, and procedures in ichnology. New York, NY: Springer-Verlag. 562 pp.
- Garrison, E.G., C.P. Giammona, F.J. Kelly, A.R. Tripp, and G.A. Wolf. 1989. Historic shipwrecks and magnetic anomalies of the northern Gulf of Mexico: Reevaluation of archaeological resource management zone 1. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 89-0024. 241 pp.
- Geraci, J.R. and D.J. St. Aubin. 1980. Offshore petroleum resource development and marine mammals. A review and research recommendations. *Marine fisheries review*, 42:1-12.
- Gordon, J.C.D., D. Gillespie, L.E. Rendell, and R. Leaper. 1996. Draft report on playback of ATOC like sounds to sperm whales (*Pheseter macrocephalus*) off the Azores. Wildlife Conservation Research Unit, Dept. of Zoology, Oxford, UK.
- Gore, R.H. 1992. The Gulf of Mexico. Pineapple Press, Florida.
- Greene, C.R. 1986. Underwater sounds from the semisubmersible drill rig SEDCO 708 drilling in the Aleutian Islands. Sect. 1 *In: API Publ 4438*. Am. Petrol. Inst., Washington, DC. 69p.
- Harvey, J.T. and M.E. Dahlheim. 1994. Cetaceans in oil. *In: Marine Mammals and the Exxon Valdez*. T.R. Loughlin, ed. Academic Press, San Diego: 257-264.
- Hayes, S.A., E. Josephson, K. Maze-Foley, and P.E. Rosel, editors. 2018. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments -- 2017. NOAA Tech Memo NMFS-NE 245; 371 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at: <https://repository.library.noaa.gov/view/noaa/22730>. Accessed September 23, 2020.
- Hayes, S.A., E. Josephson, K. Maze-Foley, P.E. Rosel, B. Byrd, S. Chavez-Rosales, T.V.N. Cole, L. Engleby, L.P. Garrison, J. Hatch, A. Henry, S.C. Horstman, J. Litz, M.C. Lyssikatos, K.D. Mullin, C. Orphanides, R.M. Pace, D.L. Palka, M. Soldevilla, and F.W. Wenzel. 2018. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 2017. NOAA Tech Memo 245 NMFS NE-245; 371 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at: <https://repository.library.noaa.gov/view/noaa/22730>. Accessed September 23, 2020.
- Hayes S.A., E. Josephson, K. Maze-Foley, P.E. Rosel, B. Byrd, S. Chavez-Rosales, T.V.N. Cole, L.P. Garrison, J. Hatch, A. Henry, S.C. Horstman, J. Litz, MC. Lyssikatos, K.D. Mullin, C. Orphanides, R.M. Pace, D.L. Palka, J. Powell, and F.W. Wenzel. 2019. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments - 2018. NOAA Tech Memo NMFS-NE 258; 291 p. Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at: <https://repository.library.noaa.gov/view/noaa/20611>. Accessed September 23, 2020.
- Hayes, S.A., E. Josephson, K. Maze-Foley, P.E. Rosel, B. Byrd, S. Chavez-Rosales, T.V.N. Cole, L.P. Garrison, J. Hatch, A. Henry, S.C. Horstman, J. Litz, M.C. Lyssikatos, K.D. Mullin, C. Orphanides,

- R.M. Pace, D.L. Palka, J. Powell, and F.W. Wenzel. 2020. US Atlantic and Gulf of Mexico Marine Mammal Stock Assessments -- 2019. NOAA Tech Memo NMFS-NE 264; Available from: National Marine Fisheries Service, 166 Water Street, Woods Hole, MA 02543-1026, or online at: <https://www.fisheries.noaa.gov/webdam/download/109188360>. Accessed September 23, 2020.
- Hazel, J., I. R. Lawler, H. Marsh, and S. Robson. 2007. Vessel speed increases collision risk for the green turtle *Chelonia mydas*. *Endangered Species Research* 3:105-113.
- Hernandez, F.J., S. Powers, and W. Graham. 2010. Seasonal variability in ichthyoplankton abundance and seasonal composition in the northern Gulf of Mexico off Alabama. *Fish Bull* 108:193-207.
- Inoue, M., S.E. Welsh, L.J. Rouse, Jr., and E. Weeks. 2008. Deepwater currents in the eastern Gulf of Mexico: Observations at 25.5°N and 87°W. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2008-001. 95 pp. <https://espis.boem.gov/final%20reports/4305.pdf> \_ Accessed September 23, 2020.
- Jefferson, T.A., M.A. Webber, and R.L. Pitman. 2008. *Marine Mammals of the World: A Comprehensive Guide to Their Identification*. Amsterdam: Elsevier. 573 pp.
- Jensen, A. S., and G. K. Silber. 2004. Large Whale Ship Strike Database. U.S. Department of Commerce, NMFS-OPR-25.
- Johansen, O., H. Rye, A.G. Melbye, H.V. Jensen, B. Serigstad, and T. Knutsen. 2001. Deep Spill JIP – Experimental discharges of gas and oil at Helland Hansen – June 2000: Final technical report. 159 pp.
- Kennicutt II, M.C., ed. 1995. Gulf of Mexico offshore operations monitoring experiment, Phase I: Sublethal responses to contaminant exposure, final report. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 95-0045. 709 pp.
- Kennicutt II, M.C., R.H. Green, P. Montagna, and P.F. Roscigno. 1996. Gulf of Mexico offshore operations monitoring experiment (GOOMEX), Phase I: Sublethal responses to contaminant exposure—introduction and overview. *Can. J. Fish. Aquat. Sci.* 53:2540-2533.
- Kessler, J.D., D.L. Valentine, M.C. Redmond, M. Du., E.W. Chan, S.D. Mendes, E.W. Quiroz, C.J. Villanueva, S.S. Shusta, L.M. Werra, S.A. Yvon-Lewis, and T.C. Weber. 2011. A persistent oxygen anomaly reveals the fate of spilled methane in the deep Gulf of Mexico. *Science Express*, 10.1126/science.1199697. Internet website: <https://science.sciencemag.org/content/331/6015/312>. Accessed September 23, 2020.
- Krivor, M.C., J. de Bry, N.J. Linville, and D.J. Wells. 2011. Archival investigations for potential colonial-era shipwrecks in ultra-deep water within the Gulf of Mexico. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Regulation and Enforcement, Gulf of Mexico OCS Region, New Orleans, LA. 158 pp.
- Laist, D.W., A.R. Knowlton, J.G. Mead, A.S. Collet, and M. Podesta. 2001. Collisions between ships and whales. *Marine Mammal Science* 17(1):35-75.
- Lugo-Fernández, A., D.A. Ball, M. Gravois, C. Horrell, and J.B. Irión. 2007. Analysis of the Gulf of Mexico's Veracruz-Havana route of *La Flota de la Nueva España*. *Journal of Maritime Archaeology* 2:24-47. July 2007.
- Lutcavage, M., P. Plotkin, B. Witherington, and P.L. Lutz. 1997. Human impacts on sea turtle survival. In: Lutz, P.L. and J.A. Musick, eds. *The biology of sea turtles*. Boca Raton, FL: CRC Press. Pp. 387-409.
- Lutz, P.L. and M. Lutcavage. 1989. The effects of petroleum on sea turtles: Applicability to Kemp's ridley. In: Caillouet, C.W., Jr. and A.M. Landry, Jr., eds. *Proceedings of the First International Symposium on Kemp's Ridley Sea Turtle Biology, Conservation and Management*. Texas A&M University Sea Grant College Program, Galveston. TAMU-SG-89-105. Pp. 52-54.

- MacDonald, I.R., ed. 1992. Chemosynthetic ecosystems study literature review and data synthesis: Volumes I-III. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 92-0033 through 92-0035.
- MacDonald, C.P., P.T. Roberts, M.R. Lilly, C.A. Knoderer, and D.S. Weeks. 2004. Boundary Layer Study in the Western and Central Gulf of Mexico. Final Report. Prepared for the U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2004-060. 588 pp.
- Miller, M.H. and C. Klimovich. 2017. Endangered Species Act Status Review Report: Giant Manta Ray (*Manta birostris*) and Reef Manta Ray (*Manta alfredi*). NMFS.
- Moore, S.F. and R.L. Dwyer. 1974. Effects of oil on marine organisms: A critical assessment of published data. *Water Research* 8:819-827.
- Murawski, S. A., W. T. Hpgarth., E. B. Peebles, and L. Barbeiri. 2014. Prevalence of External Skin Lesions and Polycyclic Aromatic Hydrocarbon Concentrations in Gulf of Mexico Fishes, Post-Deepwater Horizon, *Transactions of the American Fisheries Society*, 143:4, 1084-1097.
- Muller-Karger, F.E., F. Vukovich, R. Leben, B. Nababan, C. Hu, and D. Myhre. 2001. Surface circulation and the transport of the Loop Current into the northeastern Gulf of Mexico: Final report. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2001-102. 39 pp.
- National Research Council (NRC). 1983. Drilling discharges in the marine environment. Panel on Assessment of Fates and Effects of Drilling Fluids and Cuttings in the Marine Environment. Marine Board, Commission on Engineering and Technical Systems, National Research Council. Washington, DC: National Academy Press.
- National Research Council (NRC). 1990. The decline of sea turtles: Causes and prevention. Washington, DC: National Academy Press. 183 pp.
- National Research Council (NRC). 2003. Oil in the sea III: Inputs, fates, and effects (Committee on Oil in the Sea: J.N. Coleman, J. Baker, C. Cooper, M. Fingas, G. Hunt, K. Kvenvolden, J. McDowell, J. Michel, K. Michel, J. Phinney, N. Rabalais, L. Roesner, and R. B. Spies). Washington, DC: National Academy Press. 265 pp.
- National Research Council (NRC). 2005. Oil Spill dispersants: Efficacy and effects. Washington, DC: National Academy Press. 377 pp.
- Neal Adams Firefighters, Inc. 1991. Joint industry program for floating vessel blowout control. Prepared for the U.S. Dept. of the Interior, Minerals Management Service. TA&R Project 150.
- Neff, J.M. 1990. Composition and fate of petroleum and spill-treating agents in the marine environment. In: Geraci, J.R. and D.J. St. Aubin, eds. *Sea mammals and oil: Confronting the risks*. San Diego, CA: Academic Press, Inc. Pp. 1-33.
- Neff, J.M., T.C. Sauer, and N. Maciolek. 1989. Fate and effects of produced water discharges in nearshore marine waters. Prepared for the American Petroleum Institute, Washington, DC.
- Oil Spill Commission. 2011a. Deepwater: The Gulf oil spill disaster and the future of offshore drilling. Report to the President (ISBN: 978-0-16-087371-3). <http://www.gpo.gov/fdsys/pkg/GPO-OILCOMMISSION/pdf/GPO-OILCOMMISSION.pdf>. Accessed September 23, 2020.
- Oil Spill Commission. 2011b. The use of surface and subsea dispersants during the BP *Deepwater Horizon* oil spill. Staff Working Paper No. 4., Updated January 11, 2011.
- PCCI Marine and Environmental Engineering (PCCI). 1999. Oil spill containment, remote sensing and tracking for deepwater blowouts: status of existing and emerging technologies. Report prepared for the U.S. Dept. of the Interior, Minerals Management Service. TA&R Project 311. 66 pp. + apps. Internet website: <https://www.bsee.gov/research-record/osrr-311-oil-spill-containment-remote-sensing-and-tracking-deep-water-blowouts>. Accessed September 23, 2020.

- Pearson, C.E., S.R. James, Jr., M.C. Krivor, and S.D. El Darragi. 2003. Refining and revising the Gulf of Mexico Outer Continental Shelf Region high-probability model for historic shipwrecks. 3 vols. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2003-060 through 2003-062.
- Popper, A.N., A.D. Hawkins, R.R. Fay, D. Mann, S. Bartol, T. Carlson, S. Coombs, W.T. Ellison, R. Gentry, M.B. Halvorsen, S. Løkkeborg, P. Rogers, B.L. Southall, D. Zeddies, and W.N. Tavolga. 2014. Sound Exposure Guidelines for Fishes and Sea Turtles: A Technical Report prepared by ANSI-Accredited Standards Committee S3/SC1 and registered with ANSI. ASA S3/SC1.4 TR-2014. Springer and ASA Press, Cham, Switzerland.
- Preen, A. 1989. Observations of mating behaviour in dugongs (*Dugong dugon*). *Marine Mammal Science* 5(4): 382-387.
- Pritchard, P.C.H. 1997. Evolution, phylogeny, and current status. In: Lutz, P.L. and J.A. Musivk, eds. *The biology of sea turtles*. Boca Raton, FL: CRC Press. Pp. 1-28.
- Rawls, J.K. and D. Bowker-Lee. 2011. Shipwreck research in the New Orleans Notarial Archives. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Regulation and Enforcement, Gulf of Mexico OCS Region, New Orleans, LA., OCS Study BOEMRE 2011-040.
- Richardson, W.J., C.R. Greene, C.I. Mame, and D.H. Thomson. 1995. *Marine mammals and noise*. San Diego, CA: Academic Press Inc.
- Ridgway, S.H., and D. Carder. 2001. "Assessing hearing and sound production in cetacean species not available for behavioral audiograms: experience with *Physeter*, *Kogia*, and *Eschrichtius*" *Aquat. Mammals* 27, 267–276.
- Rowe, G. and M.C. Kennicutt, eds. 2009. Northern Gulf of Mexico continental slope habitats and benthic ecology: Final report. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2001-091. 158 pp.
- Sadiq, M. and J.C. McCain. 1993. *The Gulf War aftermath: An environmental tragedy*. Dordrecht: Kluwer Academic Publishers.
- Schroeder, W.W., S.D. Brooke, J.B. Olson, B. Phaneuf, J.J. McDonough III, and P. Etnoyer. 2005. Occurrence of deep-water *Lophelia pertusa* and *Madrepora oculata* in the Gulf of Mexico. In: Freiwald, A. and J. Murray Roberts, eds. *Cold-Water Corals and Ecosystems*. New York, NY: Springer Berlin Heidelberg. 1,243 pp.
- Shaver, D.J., K.M. Hart, I. Fujisaki, C. Rubio, A.R. Sartain, J. Peña, P.M. Burchfield, D. Gomez Gamez, and J. Ortiz. 2014. Foraging area fidelity for Kemp's ridleys in the Gulf of Mexico.
- Southall, B.L., A.E. Bowles, W.T. Ellison, J.J. Finneran, R.L. Gentry, C.R. Greene, Jr., D. Kastak, D.R. Ketten, J.H. Miller, P.E. Nachtigall, W.J. Richardson, J.A. Thomas, and P. Tyack. 2007. *Marine Mammal Noise Exposure Criteria: Initial Scientific Recommendations*. *Aquatic Mammals* 33: 411-521.
- State of Florida. Fish and Wildlife Conservation Commission. 2020. Manatee synoptic surveys. Internet website: <http://myfwc.com/research/manatee/research/population-monitoring/synoptic-surveys/>. Accessed September 23, 2020.
- St. Aubin, D.J. and V. Lounsbury. 1990. "Oil Effects on Manatees: Evaluating the Risks," Chapter 11: J.R. Geraci and D.J. St. Aubin (eds.), *Sea Mammals and Oil: Confronting the Risks*. San Diego, California: Academic Press, Inc., pp. 241-251.
- Sulak, K.J., M.T. Randall, K.E. Luke, A.D. Norem, and J.M. Miller, eds. 2008. Characterization of northern Gulf of Mexico deepwater hard-bottom communities with emphasis on *Lophelia* coral—*Lophelia* reef megafaunal community structure, biotopes, genetics, microbial ecology, and geology. U.S. Dept. of the Interior, Geological Survey (USGS Open-File Report 2008-1148) and Minerals Management Service, Herndon, VA (OCS Study MMS 2008-015).

- Teas, W.G. 1994. Marine turtle stranding trends, 1986-1993. In: Bjorndal, K.A., A.B. Bolten, D.A. Johnson, and P.J. Eliazar, compilers. Proceedings of the Fourteenth Annual Symposium on Sea Turtle Biology and Conservation. NOAA Tech. Memo. NMFS-SEFSC-351. Pp. 293-295.
- Turner, R.E., N.N. Rabalais, E.M. Swenson, M. Kasprzak, and T. Romaine. 2005. Summer hypoxia in the northern Gulf of Mexico and its prediction from 1978 to 1995. *Marine Environmental Research*, 59, 65-77.
- U.S. Army Corps of Engineers New Orleans District: The Mississippi River Drainage Basin. 2020. <http://www.mvn.usace.army.mil/Missions/MississippiRiverFloodControl/MississippiRiverTributaries/MississippiDrainageBasin.aspx>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS). 2007. Endangered Species Act Section 7 consultation on the effects of the five-year outer continental shelf oil and gas leasing program (2007-2012) in the Central and Western Planning Areas of the Gulf of Mexico. Biological Opinion. June 29. F/SER/2006/02611. 127 pp.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS). 2020a. Biological Opinion of the Federally Regulated Oil and Gas Program Activities in the Gulf of Mexico and Appendices. Internet website: <https://www.fisheries.noaa.gov/resource/document/biological-opinion-federally-regulated-oil-and-gas-program-activities-gulf-mexico>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS). 2020b. 2018-2020 Bottlenose Dolphins Unusual Mortality Event Southwest Florida. Internet website: <https://www.fisheries.noaa.gov/southeast/marine-life-distress/2018-2020-bottlenose-dolphin-unusual-mortality-event-southwest>. Last updated March 6, 2020. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS) and U.S. Dept. of the Interior (USDOI), Fish and Wildlife Service (FWS). 2007a. Green sea turtle (*Chelonia mydas*); 5-year review: Summary and evaluation. U.S. Dept. of Commerce, National Marine Fisheries Service, Office of protected Resources, Silver Spring, MD, and U.S. Dept. of the Interior, Fish and Wildlife Service, Southeast Region, Jacksonville Ecological Service Field Office, Jacksonville, FL. 102 pp. Internet website <http://www.fws.gov/northflorida/SeaTurtles/2007-Reviews/2007-green-turtle-5-year-review-final.pdf>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS) and U.S. Dept. of the Interior (USDOI), Fish and Wildlife Service (FWS). 2007b. Kemp's ridley sea turtle (*Lepidochelys kempii*); 5-year review: Summary and evaluation. U.S. Dept. of Commerce, National Marine Fisheries Service, Office of protected Resources, Silver Spring, MD, and U.S. Dept. of the Interior, Fish and Wildlife Service, Southeast Region, Jacksonville Ecological Service Field Office, Jacksonville, FL. 50 pp. Internet website: <http://www.fws.gov/northflorida/SeaTurtles/2007-Reviews/2007-Kemps-ridley-turtle-5-year-review-final.pdf>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS) and U.S. Dept. of the Interior (USDOI), Fish and Wildlife Service (FWS). 2007c. Loggerhead sea turtle (*Caretta caretta*); 5-year review: Summary and evaluation. U.S. Dept. of Commerce, National Marine Fisheries Service, Office of protected Resources, Silver Spring, MD, and U.S. Dept. of the Interior, Fish and Wildlife Service, Southeast Region, Jacksonville Ecological Service Field Office, Jacksonville, FL. 65 pp. Internet website: <http://www.fws.gov/northflorida/SeaTurtles/2007-Reviews/2007-Loggerhead-turtle-5-year-review-final.pdf>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS) and U.S. Dept. of the Interior, Fish and Wildlife Service (FWS). 2007d. Leatherback sea turtle (*Caretta caretta*); 5-year review: Summary and evaluation. U.S. Dept. of Commerce, National Marine Fisheries Service, Office of protected Resources, Silver Spring, MD, and U.S. Dept. of the Interior, Fish and Wildlife Service, Southeast Region, Jacksonville Ecological Service Field Office, Jacksonville, FL. 65 pp. Internet website: <http://www.fws.gov/northflorida/SeaTurtles/2007-Reviews/2007-leatherback-turtle-5-year-review-final.pdf>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS) and U.S. Dept. of the Interior, Fish and Wildlife Service (FWS). 2007e. Hawksbill sea turtle (*Eretmochelys imbricata*); 5-

- year review: Summary and evaluation. U.S. Dept. of Commerce, National Marine Fisheries Service, Silver Spring, MD, and U.S. Dept. of the Interior, Fish and Wildlife Service, Southeast Region, Jacksonville Ecological Service Field Office, Jacksonville, FL. 90 pp. Internet website: Accessed May 26, 2015. <http://www.fws.gov/northflorida/SeaTurtles/2007-Reviews/2007-hawksbill-turtle-5-year-review-final.pdf>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS) and U.S. Dept. of the Interior (USDOI), Fish and Wildlife Service (FWS). 2013a. Hawksbill sea turtle (*Eretmochelys imbricata*); 5-year review: Summary and evaluation. U.S. Dept. of Commerce, National Marine Fisheries Service, Office of protected Resources, Silver Spring, MD, and U.S. Dept. of the Interior, Fish and Wildlife Service, Southeast Region, Jacksonville Ecological Service Field Office, Jacksonville, FL. 92 pp. Internet website: <https://repository.library.noaa.gov/view/noaa/17041>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Marine Fisheries Service (NMFS) and U.S. Dept. of the Interior (USDOI), Fish and Wildlife Service (FWS). 2013b. Leatherback sea turtle (*Dermochelys coriacea*); 5-year review: Summary and evaluation. U.S. Dept. of Commerce, National Marine Fisheries Service, Office of protected Resources, Silver Spring, MD, and U.S. Dept. of the Interior, Fish and Wildlife Service, Southeast Region, Jacksonville Ecological Service Field Office, Jacksonville, FL. 92 pp. Internet website: <https://repository.library.noaa.gov/view/noaa/17029>. Accessed September 23, 2020.
- U.S. Dept. of Commerce (USDOC), National Oceanic and Atmospheric Administration (NOAA). 2010. *Deepwater Horizon* oil spill: Characteristics and concerns. Internet website: [https://web.archive.org/web/20170427015646/http://sero.nmfs.noaa.gov/deepwater\\_horizon/documents/pdfs/fact\\_sheets/oil\\_characteristics.pdf](https://web.archive.org/web/20170427015646/http://sero.nmfs.noaa.gov/deepwater_horizon/documents/pdfs/fact_sheets/oil_characteristics.pdf). Accessed September 23, 2020.
- U.S. Dept. of the Interior (USDOI), Bureau of Ocean Energy Management (BOEM). 2012. Gulf of Mexico OCS oil and gas lease sales; 2012-2017; Western Planning Area Lease Sales 229, 233, 238, 246, and 248; Central Planning Area Lease Sales 227, 231, 235, 241, and 247. final environmental impact statement. 3 vols. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Gulf of Mexico OCS Region, New Orleans, LA. OCS EIS/EIA BOEM 2012-019. [http://www.boem.gov/Environmental-Stewardship/Environmental-Assessment/NEPA/BOEM-2012-019\\_v1.aspx](http://www.boem.gov/Environmental-Stewardship/Environmental-Assessment/NEPA/BOEM-2012-019_v1.aspx). Accessed September 23, 2020.
- U.S. Dept. of the Interior (USDOI), Bureau of Ocean Energy Management (BOEM). 2017a. Gulf of Mexico OCS Oil and Gas Lease Sales: 2017-2022; Gulf of Mexico Lease Sales 249, 250, 251, 252, 253, 254, 256, 257, 259, and 261-Final Environmental Impact Statement (Multisale EIS). 3 vols. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Gulf of Mexico OCS Region, New Orleans, LA. OCS EIS/EA BOEM 2017-009. Internet website: <https://www.boem.gov/BOEM-EIS-2017-009-v1/>. Accessed September 23, 2020.
- U.S. Dept. of the Interior (USDOI), Bureau of Ocean Energy Management (BOEM). 2017b. Gulf of Mexico OCS Lease Sale Final Supplemental Environmental Impact Statement 2018 (SEIS 2018). 2 vols. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Gulf of Mexico OCS Region, New Orleans, LA. OCS EIS/EA BOEM 2017-074. Internet website: <https://www.boem.gov/BOEM-EIS-2017-074-v1/>. Accessed September 23, 2020.
- U.S. Dept. of the Interior (USDOI), Bureau of Ocean Energy Management (BOEM). 2017c. Catastrophic Spill Event Analysis: High-volume, extended-duration oil spill resulting from loss of well control on the Gulf of Mexico outer continental shelf; 1<sup>st</sup> revision. U.S. Dept. of the Interior, Bureau of Ocean Energy Management, Gulf of Mexico OCS Region, New Orleans, LA. OCS Report BOEM 2017-007.
- U.S. Dept. of the Interior (USDOI), Bureau of Safety and Environmental Enforcement (BSEE). 2020. Incident Statistics and Summaries. Internet website: <http://www.bsee.gov/Inspection-and-Enforcement/Accidents-and-Incidents/Listing-and-Status-of-Accident-Investigations/>. Accessed September 8, 2020.
- U.S. Department of Interior (USDOI), Fish and Wildlife Service (FWS). 2018. Biological Opinion Oil and Gas Leasing, Exploration, Development, Production, Decommissioning, and All Related Activities in the Gulf of Mexico Outer Continental Shelf. Issued April 20, 2018.

- U.S. Dept. of the Interior (USDOI), Minerals Management Service (MMS). 2000. Gulf of Mexico deepwater operations and activities: Environmental assessment. U.S. Dept. of the Interior, Minerals Management Service.
- U.S. Dept. of the Transportation (USDOT), Federal Aviation Administration (FAA). 2004. Visual Flight Rules (VFR) Flight Near Noise-Sensitive Areas. Advisory Circular No. 91-36D. 2 pp. Internet website: [https://www.faa.gov/documentLibrary/media/Advisory\\_Circular/AC\\_91-36D.pdf](https://www.faa.gov/documentLibrary/media/Advisory_Circular/AC_91-36D.pdf) Accessed September 23, 2020.
- U.S. Environmental Protection Agency (USEPA). 2020. Nonattainment Areas for Criteria Pollutants (Green Book). Last updated August 31, 2020. Internet Website: <https://www.epa.gov/green-book>. Accessed September 23, 2020.
- Vanderlaan, A.S., and C.T. Taggart. 2007. Vessel collisions with whales: The probability of lethal injury based on vessel speed. *Marine Mammal Science* 23(1):144-156.
- Vanderlaan, A.S.M., J.J. Corbett, S.L. Green, J.A. Callahan, C. Wang, R.D. Kenney, C.T. Taggart, and J. Firestone. 2009. Probability and mitigation of vessel encounters with North Atlantic right whales. *Endangered Species Research* 6:273-285.
- Wartzok, D., and D.R. Ketten. 1999. Marine mammal sensory systems. In *Biology of Marine Mammals* (Ed. by J. E. Reynolds and S. A. Rommel), pp. 117-175. Washington, D.C.: Smithsonian Institution Press.
- Wells, R.S. and M.D. Scott. 1997. Seasonal incidence of boat strikes on bottlenose dolphins near Sarasota, Florida. *Marine Mammal Science* 13:475-480. Wells, R.S. and M.D. Scott. 1997. Seasonal incidence of boat strikes on bottlenose dolphins near Sarasota, Florida. *Marine Mammal Science* 13:475-480.
- Wells, R.S. and M.D. Scott. 1997. Seasonal incidence of boat strikes on bottlenose dolphins near Sarasota, Florida. *Marine Mammal Science* 13:475-480.
- Welsh, S.E., M. Inoue, L.J. Rouse, Jr., and E. Weeks. 2009. Observation of the deepwater manifestation of the Loop Current and Loop Current rings in the eastern Gulf of Mexico. U.S. Dept. of the Interior, Minerals Management Service, Gulf of Mexico OCS Region, New Orleans, LA. OCS Study MMS 2009-050. 110 pp.
- Würsig, B., T. Jefferson, and D. Schmidly. 2000. The marine mammals of the Gulf of Mexico. College Station, TX: Texas A&M University Press.

## 7. PREPARERS

### NEPA Coordinator

Andrea Heckman

Environmental Scientist

### Contributors

Stacie Merritt

Tre Glenn

Physical Scientist—Air Quality Issues

Biologist—Marine Mammals, Sea Turtles, Protected Species Issues

Cholena Ren

Arie Kaller

Physical Scientist—Water Quality Issues

Biologist—Essential Fish Habitat, Commercial Fishing and Fish Resources

Jeri Wisman

Doug Jones

Biologist—Benthic Communities

Marine Archaeologist—Archaeological Issues

### Reviewers

Annette Ehrhorn

Perry Boudreaux

Senior Environmental Scientist

Unit Chief, Environmental Operations Section

## 8. APPENDIX

Appendix A—Accidental Oil-Spill Discussion

Available: <http://www.boem.gov/Appendix-A-Deepwater-SEA-Oil-Spill-Discussion/>